

SUMMARY REPORT 2012



**SHINING TREE GOLD PROJECT,
SHINING TREE ONTARIO, CANADA**

Signing Date: June 18, 2012

James R. Trusler, P.Eng



SHINING TREE SUMMARY REPORT

EXECUTIVE SUMMARY

Platinex's Shining Tree gold project comprises 50 claims (139 claim units) and one mining lease for a total of 2,240 ha (5,600 acres) and is located in the Larder Lake Mining Division, Ontario. Highway 560 midway between Sudbury and Timmins, bisects the property. The claims are held as to 100% by Platinex subject to a 3% NSR to Skead Holdings and a 100% interest in the mining lease with no underlying interests. Five of the above claims containing eleven claim units and four additional claims comprising ten claim units held by Creso Resources Inc. are part of the Gold Corona joint venture. The joint venture is not the subject of this report

The Company SEDAR-filed its NI 43-101 technical report "the Report" dated October 2, 2008, by J.G. Bryant and D. Jamieson which examines the Herrick gold deposit on Shining Tree property and seven other known gold prospects, some of which have been explored underground. The Report qualifies the sampling and drilling work by Unocal (1989) and Fort Knox (1990) compliant to NI 43-101 standards. This report draws on the previous Report and provides an update based on results of subsequent exploration.

The Report states that, "Judging from evidence of previous results on the Herrick deposit there is a potential to outline a small commercial gold deposit." An internal valuation of the Herrick deposit completed in May, 2012 by the author estimated potential gold content ranging from 111,000 oz to 172,000 oz Au within the defined vein structures to a depth of 300m. The estimate is based on a minimum of 1,628,000 tonnes grading 2.11g/tonne to a maximum of 2,554,000 tonnes grading 2.10g/tonne Au. The estimate is derived from 3 narrow sub-parallel quartz vein breccias and an enclosing mineralized shear zone within a 370m strike length, to a 300m depth and within an 80m wide block. A cut off grade of 0.5g/tonne was used in the calculations.

The Central zone of the Herrick deposit outcrops for most of its length and is a continuously readily definable body. The valuation also identified thicker gold zones between intersecting or bifurcating veins in some holes (e.g. hole HP10-44 to the southwest) and in outcrops which are not included in the calculations. These structures plunge near vertically providing a prospective target for deeper exploration. There is an undefined potential for a southwesterly plunging thickened enriched section that could contain from a depth of 300m to 1500m some 14.6 million tonnes at a grade ranging from 2.0 to 5.0 g/tonne Au for a gold content of 1.0 to 2.4 million ounces. The model is based on the mentioned structure, comparison to the West Timmins deposit and statistical grade estimates by the author. The potential quantity and grade is conceptual in nature. There has been insufficient exploration to define a mineral resource and it is uncertain if further exploration will result in the delineation of a mineral resource. The internal valuation was conducted by the author, who is a qualified person but an insider of the issuer and therefore according to the ethical standards set forth in the National Instrument 43-101 no qualification of resources is permitted. More sampling of existing core and stripping of identified thicker mineralized zones where these project to surface, is warranted and could increase the identified potential within the exploration envelope and ultimately lead to qualification of resources by an independent qualified person.

Since October 2008, the Company has carried out an exploration program on the property. Particular interest is focused on 446 samples recovered from basal till that have been processed by Overburden Drilling Management. The results have revealed one of the more exceptional gold dispersion trains in the Abitibi. Many of the samples contain pristine gold grains indicating a nearby bedrock source. In addition, 169 samples each returned more than ten gold grains (up to 144 gold grains in one sample) that suggest several nearby previously unexplored gold occurrences in the bedrock. Noticeably a line

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

of the highest counts was obtained along 9 km easterly trending shear believed to be the Larder Lake Break or an associated splay fault and correlating with broad XDS ORTHO anomalies, a potassium decay anomaly, circular magnetic anomalies, and partly circumscribing and correlating with an alkalic stock. The occurrence, distribution and density of high gold counts in till is similar to such patterns evident close to major gold deposits throughout the Timmins and Matachewan gold camps. The pattern encourages the conclusion that the sources of such anomalous indications are expressions of one or more major gold deposits.

A nine km Pole-dipole IP survey was completed in March, 2012 by JVX Ltd. This was immediately followed by a five hole 870m drilling program, also in March, 2012, on previously undrilled portions of the property to test five very strong IP anomalies spatially associated with gold in till anomalies.

All of the holes intersected thick zones of mineralization and/or alteration. Logging, sampling and assaying of the core is partly completed and initial results reported. One hole intersected unmineralized, chlorite altered, syenite with a few randomly selected samples returning anomalous values to 327ppb Au. The strong IP response was unexplained. This area and its strike projections are a high priority for follow up exploration. Furthermore the evident clustering of the higher gold in till counts near the margins of an alkalic or syenitic stock is suggestive of a prominent, recognizable and readily explored, geological environment on a large portion of the property.

On a regional scale the property is located within the prolific Abitibi greenstone belt which is renowned for its concentration of world-class gold and Cu-Zn-Ag-Au deposits. The Shining Tree property appears to be on trend with the Larder Lake – Cadillac Break with which many of the world-class gold deposits are spatially associated. The Shining Tree property is also centred on the axis of a regional gravity high signifying an area of thick supracrustal rocks and under-plating by oceanic crust or an intrusive environment. In the Abitibi these associations appear conducive to the genesis of major gold deposits.

The property is underlain by an Early Precambrian assemblage of WNW trending, steeply dipping felsic to mafic and ultramafic metavolcanic rocks as well as metasedimentary rocks and alkali metavolcanic rocks of Timiskaming age. These rocks are intruded by pre-orogenic feldspar porphyries, alkali (syenite) stocks and hypabyssal diabase dikes of Matachewan age and sills of Nipissing age. The Timiskaming-aged metasedimentary rocks are believed to have accumulated in a regional rift. Later south to north compressive faulting along the axis of this rift is proposed, but has not been documented in the Shining Tree area. Major NNW sinistral wrench faults, including the Michiwakenda Lake Fault which crosses the Shining Tree property, are regional in scale. Related subsidiary parallel shears and possible splays carry significant gold mineralization on the Shining Tree property.

Bedrock exposures are limited on the property which is principally covered by a thin mantle of Wisconsinan till.

The Platinex Shining Tree property is at the hub of the area known as the Shining Tree gold camp. Numerous old gold workings and showings are located along a north-northwest trend on the property and to the southeast of the property for exploration. There are other workings and showings in one or two parallel trends. Minor production has been achieved from one set of workings close to the property boundary.

Gold-silver mineralization is principally associated with multiple brittle, dilatant (but sometimes ductile) quartz-carbonate veins. Many of these veins are narrow and contain sporadic high gold (silver) values, but are of lower priority for exploration. The Herrick gold deposit discovered in 1918 has been developed by a shaft to 94 metres and some 345 m of lateral development on two veins. Very high grade shoots were defined in surface channel sampling, sampling in the shaft, underground on the level at 90 m and in four drill holes by Herrick Gold Mines from 1918 onwards. In 1989 Unocal drilled 11

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

holes for 1,473 m, cut over 201 channel samples and also took 35 composite grab samples on the two veins to test a stated potential, solely from historic data, of the system to carry 1,732 tonnes per vertical foot at 7.2 g Au /t over a width of 1.8 m and 50% dilution. Unocal's results confirmed the persistence of the gold mineralization over a 385 metre strike length and expanded the known thickness of the mineralized shear along much of its length. However, the drilling and channel sampling by Unocal identified that certain gold-bearing zones are narrowly confined within the shear zones and carry potentially commercial, high grade, correlative values and other portions contain isolated patches of high gold grades within a broader lower grade shear zone.

Although the tenor of the Herrick Gold Mines drill core and channel sampling was not confirmed completely, the location of the mineralization was confirmed by Unocal. Surface sample gold values of up to 56.5g per tonne /1.0 m (45 samples) were reported. In 1990 Fort Knox Mines performed a small program of channel sampling on surface and in the shaft confirming some of the higher grade mineralization but not all of the higher grade values presented by Herrick Gold Mines.

To-date work by Platinex has included: an airborne magnetometer, VLF-EM and radiometric survey; stripping, channel sampling (58 samples on Herrick) and core sampling conducted to verify results of previous work on the Herrick, Caswell and Churchill workings; a detailed program of basal till sampling collecting 446 samples now processed for gold grains by Overburden Drilling Management; prospecting, sampling and geological mapping; 98.8 line km ground walking magnetometer surveys and 12km IP surveys; 51 drill holes for 6,181m on the Herrick deposit, 7 holes for 1,070m on the Caswell area, and 5 holes for 870m to test coincident IP and gold in till anomalies. Further whole rock analyses of the fines fraction was carried out on 205 till samples. Most parameters can be related to nearby bedrock features which provides additional evidence of the proximal source of the gold grains.

One conclusion that can be made from the analyses of fines fractions of basal till samples and drilling to date is that the East West shears found at the Caswell prospect display a gold-silver tellurium association throughout the extensive set of trenches whereas the Michiwakenda Lake fault and other NNW trending structures host a gold-arsenic association. The relationship between the two systems is unknown. However it is postulated that the alkalic stock found in the middle of the claim group may be the driving heat source for the other peripheral deposits and that the emphasis on exploration should be directed to those anomalies associated with the stockf and its periphery.

A two-phase exploration program is proposed to step up the evaluation of existing and anticipated gold-bearing targets during the balance of 2012 and in 2013. The proposed work program should include more core sampling, stripping and channel sampling, then deeper diamond drilling of the gold zones at the Herrick deposit and testing for the depth continuity of the Ronda gold-bearing zone beneath the Platinex property. Additional detailed surface prospecting, surface stripping and channel sampling, line cutting, pole-dipole IP, soil gas surveys and diamond drilling are also recommended. The proposed budget for Phase I is \$1,554,500, and the budget for Phase II is \$3,272,550.

Proposed drilling in Phase 1 totals 5,000 m and in Phase II 15,000m.

Table of Contents

EXECUTIVE SUMMARY	ii
1.0 INTRODUCTION AND GENERAL INFORMATION	10
1.1 Introduction.....	10
1.2 Terms of Reference.....	10
1.3 Scope of Work and Acknowledgements.....	10
1.4 Sources of Information	11
1.5 Disclaimer.....	11
2.0 PROPERTY LOCATION, ACCESS, DESCRIPTION AND OWNERSHIP	11
2.1 Property Location and Access.....	11
2.2 Property Description.....	12
2.3 Property Ownership.....	12
2.4 Access, Local Resources and Infrastructure.....	12
3.0 PHYSIOGRAPHY, TOPOGRAPHY, CLIMATE AND ENVIRONMENT	13
3.1 Physiography and Topography	13
3.2 Climate	13
3.3 Land Uses, Environmental and Safety Considerations	13
4.0 COMMUNITIES AND FIRST NATIONS	14
4.1 Communities	14
4.2 First Nations	14
5.0 GEOLOGY	14
5.1 Quaternary Geology	14
5.2 Regional Geology and Bedrock Assemblages	15
5.2.1 Assemblages	15
5.3 Structural Geology.....	17
6.0 MINERAL DEPOSIT TARGETS	18
6.1 Gold Deposit Model.....	18

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

6.1.1	Gold Potential of Platinex Shining Tree Property	19
6.2	Volcanogenic Massive Sulphide Deposits (VMS) Model.....	20
6.2.1	VMS Potential of Platinex Shining Tree Property	20
7.0	DEPOSITS AND OCCURRENCES OF THE SHINING TREE AREA PROPERTY AND LOCAL AREA GEOLOGY	21
7.1	Past Producers.....	21
7.1.1	Ronda Mine	21
7.1.2	Tyranite Mine	21
7.1.3	Bilmac Mine	21
7.2	Occurrences with Past Underground Development	21
7.2.1	Herrick Deposit	22
7.2.2	Caswell Occurrence.....	23
7.2.3	Churchill Occurrence	23
7.2.4	Other Underground Development Sites.....	24
7.3	Occurrences and Prospects	24
7.3.1	On-Property	24
7.3.2	Occurrences in the General Area	24
8.0	PROPERTY GEOLOGY	25
8.1	Herrick Deposit.....	25
8.2	Caswell Area	27
8.3	Churchill Area.....	28
8.4	Speed Lake area	29
9.0	MINERALIZATION.....	30
9.1	Herrick	30
9.2	Caswell area.....	31
9.3	Speed Lake area	31
10.0	EVALUATION OF SHOWINGS	31
10.1	Caswell area.....	31

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

10.2	Speed Lake Area.....	33
10.3	Beilby Lake Area	34
11.0	GEOPHYSICAL SURVEYS	34
11.1	Regional Gravity Surveys.....	34
11.2	Airborne Geophysical Surveys	35
11.2.1	Prior Surveys	35
11.3	Ground Geophysical Surveys.....	37
12.0	GEOCHEMICAL INVESTIGATIONS	37
12.1	Regional Lake Sediment Survey	37
12.2	Lithogeochemical Studies	38
12.3	On-Site Geochemical Surveys	38
12.3.1	Basal Till Sampling	38
13.0	DRILLING	40
13.1	Historic Drilling	40
13.2	Platinex Drilling.....	40
14.0	DATA COMPILATION AND SYNTHESIS.....	40
14.1	Herrick Deposit.....	40
14.2	Caswell area.....	42
14.3	Churchill area	42
14.4	Speed Lake Area.....	42
15.0	SAMPLE COLLECTION, HANDLING AND PREPARATION.....	43
16.0	SAMPLE ANALYSIS, QUALITY CONTROL, AND QUALITY ASSURANCE.....	44
17.0	DATA VALIDATION - 2008.....	45
17.1	Site Visit and Check Sampling	45
17.1.1	Herrick Deposit	45
18.0	COMPETITOR ACTIVITY.....	47
19.0	MINERAL PROCESSING AND METALLURGICAL TESTING	48

20.0 MINERAL RESOURCE AND MINERAL RESERVE ESTIMATES.....	48
21.0 DISCUSSION.....	49
21.1 Historic Context	49
21.2 Spatial Context.....	49
21.3 Caswell area.....	50
21.4 Herrick Deposit.....	50
21.5 Churchill area	51
21.6 Ribble vein (Ronda).....	51
22.0 CONCLUSIONS.....	51
23.0 RECOMMENDATIONS.....	52
24.0 REFERENCES	56
25.0 CERTIFICATES OF QUALIFICATIONS	61

LIST OF TABLES

TABLE 1: CLAIMS AND STATUS AS OF SEPTEMBER 2008
TABLE 2: TERMS AND ABBREVIATIONS
TABLE 3: VALIDATON SAMPLING RESULTS FOR THE HERRICK DEPOSIT
TABLE4: PROPOSED PHASE I BUDGET
TABLE5: PROPOSED PHASE II BUDGET
TABLE 6: PROPOSED PHASE I WORK PROGRAM

LIST OF FIGURES

FIGURE 1: GENERAL LOCATION MAP
FIGURE 2: CLAIM MAP
FIGURE 3: GEOLOGICAL SUBDIVISIONS OF ONTARIO
FIGURE 4: SIMPLIFIED GEOLOGICAL MAP OF ABITIBI GREENSTONE BELT SHOWING DISTRIBUTION OF MAJOR FAULT ZONES AND GOLD DEPOSITS
FIGURE 5: LITHOSTRATIGRAPHIC ASSEMBLAGES OF THE SHINING TREE AREA

- FIGURE 6: GEOLOGY
- FIGURE 7: HERRICK GEOLOGY AND DRILL HOLE LOCATIONS
- FIGURE 8: CASWELL PROSPECT HIGHLIGHTS-
- FIGURE 9: SKETCH OF BEILBY LAKE SHOWING
- FIGURE 10: PRELIMINARY IP SURVEY PLANS, CHARGEABILITY N=4
- FIGURE 11: AEROMAGNETIC AND ELECTROMAGNETIC SURVEY – FIRST VERTICAL DERIVATIVE MAGNETICS AND EM ANOMALIES
- FIGURE 12: AEROMAGNETIC AND ELECTROMAGNETIC SURVEY – XDS ORTHO AND EM ANOMALIES
- FIGURE 13: INTERPRETATION OF GEOPHYSICAL SURVEY RESULTS
- FIGURE 14: HERRICK GOLD DEPOSIT, CROSS SECTION 13 + 35 NORTH
- FIGURE 15: HERRICK GOLD DEPOSIT, LONGITUDINAL SECTION CENTRAL ZONE
- FIGURE 16: RELATION OF GOLD IN TILL TO MAJOR GOLD DEPOSITS
- FIGURE 17: GOLD GRAINS IN TILL
- FIGURE 18: COMPILATION: GEOLOGY, XDS ORTHO ANOMALIES WITH GOLD IN TILL
- FIGURE 19: ARSENIC IN TILL
- FIGURE 20: NICKEL IN TILL
- FIGURE 21: COPPER IN TILL
- FIGURE 22: SULPHUR IN TILL

LIST OF APPENDICES

- APPENDIX A: GLOSSARY OF TERMS AND ABBREVIATIONS
- APPENDIX B: CERTIFICATE OF CONSENT
- APPENDIX C: TABLES D1-D4; DUPLICATE AND PULP METALLIC ASSAY RESULTS FROM UNOCAL AND FORT KNOX EXPLORATION

1.0 INTRODUCTION AND GENERAL INFORMATION

1.1 Introduction

The Shining Tree property is located in the prolific Archean-age Abitibi Greenstone Belt which hosts many significant gold deposits and past producers of the Destor - Porcupine and Larder Lake - Cadillac trends in Ontario and Quebec (Figures 1 and 2). World-class volcanogenic massive sulphide deposits have also been discovered and mined over the last ninety years.

The Shining Tree area has been the focus of gold exploration since the early 1900's. There are numerous gold occurrences and minor gold production has occurred at three locations.

The property comprises a consolidation of 139 claim units by Robert Macgregor of Skead Holdings Ltd. and one mining lease acquired from Gary John McBride. Mr. Macgregor has been actively prospecting in the Abitibi Greenstone Belt for more than 40 years. Several historic gold properties, including shafts, waste and ore dumps, and mechanically stripped areas are present on the property and others are known in the Shining Tree area.

Excellent continuity of some high grade gold-bearing veins over mineable widths as at the Herrick deposit, the significant extent of previously inaccessible but now prospective overburden covered areas, the numerous occurrences of gold in the Shining Tree area, the newly identified highly prospective, glacial gold dispersion train, and the shallow nature of exploration to date all tend to provide impetus for further exploration.

Drilling by Platinex on the Shining Tree property has indicated potential for the Herrick veins to host a small scale commercial deposit or even a higher grade deposit at depth. Till sampling has provided evidence of large scale gold occurrences on the scale of a major mining camp which are yet to be explored.

1.2 Terms of Reference

James R. Trusler, P.Eng who is also President and CEO of Platinex Inc. has prepared this report as an update of information to the National Instrument 43-101 report on the property written in 2008 by J. Garry Bryant and David Jamieson. All work done since then has been supervised by independent qualified persons and a comprehensive program of exploration has been carried out on the property.

The main purpose of this summary report is to characterize the setting and objectives of the Shining Tree gold exploration project and provide a view of the potential.

1.3 Scope of Work and Acknowledgements

The objective and scope of work associated with this summary report was three-fold. First priority was to review available government publications, assessment reports filed with Ministry of Northern Development and Mines and any other reports and information owned or accessible through Skead Holdings and or Platinex Inc. The second priority was to obtain first-hand familiarization and evaluation of some of the showings and other areas of the property. Thirdly, the author has reviewed all of the field work carried out by Platinex under the supervision of David Jamieson, Scott Franko and Dean Cutting since 2008.

This summary report was prepared to meet the standards of disclosure for mineral projects with the exception that it is not independent and mineral resources cannot be included. The author utilized Canadian Securities Administrators National Instrument 43-101 (NI 43-101), Form 43-101F1 and Companion Policy 43-101CP as guidelines for report preparation. The professional opinions of Z.

Dvorak consulting geophysicist were helpful in the assessment and interpretation of the wide-spaced, government-funded AEM and magnetometer survey. He also provided advice and an interpretation regarding the airborne, fixed-wing VLF-EM, magnetometer and radiometric survey carried out on behalf of Platinex by Terraquest. Map compilation services were provided by Mohammad Salman, Jeff Meek and Associates and by Iain Trusler. Geologists working on the project since 2008 include Dave Jamieson, Garry Bryant, Mehmet Spaho, Dean Cutting, Scott Franko, Frank Racicot and Fred Gittings. Thin section Petrographic studies were provided by Fred Gittings and Richard James.

1.4 Sources of Information

The main technical documents and files related to the Shining Tree Project consist of:

- MNDM land use and land tenure files.
- OGS publications, databases and assessment reports.
- Government mapping for this area.
- Results of Overburden Drilling Management Ltd. evaluation of basal till samples.
- Drilling, prospecting, stripping, channel sampling, mapping and airborne and ground geophysical surveys supervised and/or commissioned by Platinex Inc.

1.5 Disclaimer

This report is built on the Qualifying Report prepared in 2008 which is quoted herein “The authors, including J.G. Bryant, the qualified person, attempted to confirm the validity of available information and site characteristics whenever possible. Several of the reports provided excellent evaluation and documentation of activities. Successful validation of some work was possible due to Certificates of Analysis in reports and due to independent, reasonable duplication of gold concentrations at the Herrick showing.

Confirmation of the opinions and conclusions in historic correspondence and reports prepared prior to 1989 is impossible. It was also impossible to validate most of the geological, geophysical and geochemical information included in assessment reports, exploration company websites and government publications.

All claims listed as project property correlated with claims identified as belonging to Skead Holdings Ltd. in the “Mining Claim Client Report” downloaded from the MNDM website on September 22, 2008. Very limited field validation of the claim boundaries, claim posts and claim tags has indicated that at least some boundary and post locations are as represented on Mining Recorder claim maps. The authors do not, however, warrant the validity and locations of all Platinex claims or the information provided on the MNDM claim maps.”

The claims listed in this report are claims now listed as belonging to Platinex Inc. In the Mining Claim Client Report downloaded from the MNDM website on June 7, 2012.

2.0 PROPERTY LOCATION, ACCESS, DESCRIPTION AND OWNERSHIP

2.1 Property Location and Access

The Shining Tree property is located east of Highway 144 in northeastern Ontario at a latitude about mid-way between the cities of Sudbury and Timmins (Figure 1 and 2). Highway 560 extends east from Highway 144, bypasses the former CN railway stop of Westree, passes through the hamlet of Shining Tree and then extends east to Gowganda, Elk Lake and then New Liskeard located on Highway 11.

Highway 560 traverses the property in a northeast-southwest direction and enters the south portion of the property approximately 1.7 kilometres northeast of the hamlet of Shining Tree. The property underlies portions of Macmurchy, Churchill and Asquith Townships.

The approximate centre of the property has UTM coordinates of Zone 17, 483787mE, 5271620mN. That location correlates with a longitude and latitude of 81°12'56" W and 47°35'52" N.

2.2 Property Description

The Shining Tree property comprises 46 claims totalling 139 claim units (Figure 2 and Table 1) and one mining lease. The claims comprise a larger north-south block and a smaller east-west block that have one common corner post. All claims are registered in the Larder Lake mining division. Work completed on the property to date is progressively being filed for assessment and there is currently sufficient coverage to keep the entire property in good standing for 20-25 years. Five of the claims comprising eleven claim units are subject to the Gold Corona joint venture with Creso Resources Inc.

2.3 Property Ownership

In 2011 the Company vested an option agreement with Skead Holdings Ltd., with respect to 139 claim units (5,680 acres or 2,299 ha), situated in Churchill, Macmurchy and Asquith Townships in Ontario (the "Shining Tree property"). The Company now holds a 100% interest in the claims subject to a 3% NSR and subject to an advanced annual royalty of \$10,000 commencing in April 2013. The 3% NSR may be reduced by payment of \$400,000 for each 0.5% NSR purchased to a maximum of 1.5% NSR. If the optionor wishes to sell the royalty interest the Company retains a right of first refusal to purchase the NSR. On March 30, 2012 Platinex entered into an agreement to acquire a lease (40 acres, 16 ha) from Gary John McBride for 200,000 shares of Platinex which purchase closed on May 10, 2012. The lease is central to the Shining Tree property, is renewable but set to expire on September 30, 2016.

On October 20, 2010, Platinex entered into a joint venture agreement with Creso Resources Inc. ("Creso") with the provision that a more formal agreement be produced within 3 months by Creso. That never happened but in a technical committee meeting in 2011 it was agreed that each party would be responsible for exploring its own claims. Claims belonging to Platinex are subject to the joint venture but there is no current certainty that the joint venture will be continuous. Five of Platinex's claims comprising 11 claim units subject to the joint venture include: L1227175, L1235004, L4209213, L4207972 and L4217645.

Four claims held by Creso comprising 10 claim units which also are subject to the joint venture agreement include: L1242013, L1242014, L1242015 and L4225667.

2.4 Access, Local Resources and Infrastructure

All-weather road access to the general area is excellent (Figure 2). Active lumber road and bush road access from Highway 560 to various parts of the property is fairly limited. Former roads into the north part of the main block were restored with a bridge established crossing a creek in conjunction with Creso Resources Inc. This has provided ready access to the Herrick deposit.

Some of the south part of the main block is also less accessible. Several parts of the east block are accessible via good lumber road and then lesser traveled old roads and trails.

Several tourist camps located in the area offer fully-equipped cabins for accommodation. Rental boats and motors are also available. Gasoline and a small selection of groceries and dry goods are available at the Spruce Shilling Camp. Gogama is the closest community with minor services. More substantial supplies and materials must be purchased in Timmins or Greater Sudbury.

3.0 PHYSIOGRAPHY, TOPOGRAPHY, CLIMATE AND ENVIRONMENT

3.1 Physiography and Topography

This area of northeastern Ontario is typified by extensive spruce bush mixed with some poplar and other species. The surficial geology is dominated by Quaternary ground moraines. Glaciations have created a mosaic of numerous lakes, some swamp and muskeg, several creeks and the Montreal River which flows in a northerly direction. The south part of the largest lake, Michiwakenda Lake, extends about 2.5 kilometres in a northwest-southeast direction along the northeast side of the main claim block. Other significant lakes on the west part of the property include Perkins Lake, Gosselin Lake, Cryderman Lake and Chlorus Lake. The hamlet of Shining Tree is situated a short distance to the west on the shore of Shining Tree Lake. The larger surface water bodies on the east part of the property include the southeast extremity of Michiwakenda Lake, Wasapika Lake, Knox Lake and Bob Lake (Figure 2).

The West Montreal River is the dominant drainage feature in the area. It flows north across the eastern part of the property and has a wider section called Wasapika Lake. Michiwakenda Lake, Shining Tree Creek and Caswell Lake all drain into the Montreal River from the northwest. Undulating terrain characterizes the area. Elevations range from 335 to 400 metres above mean sea level (AMSL). Highest elevations occur along a northwest-southeast trend that crosses the Perkins Lake area in the northwest part of the property and recurs in a zone that extends from a short distance northwest of Highway 560 toward the southeast beyond the property (Figure 2). There appears to be a strong correlation between at least some of the topographically higher areas and intermediate to felsic volcanic rocks and some metasedimentary rocks.

3.2 Climate

A northern temperate climate with warm summers and cold winters is characteristic for the area. The closest weather stations reported by Environment Canada (www.climate.weatheroffice.ec.gc.ca) are at Timmins, Sudbury and Earlton. Data for 1961 – 1990 indicated temperatures ranged from -45°C in winter to 40°C in summer. Average annual precipitation was 554 to 657 mm rainfall and 247 to 313 cm snow for a total of 785 to 899 mm. Maximum snow depth was 157 cm. Exploration activity can be performed year-round.

3.3 Land Uses, Environmental and Safety Considerations

The primary current activities on and in the vicinity of the claims include soft-wood lumbering operations by Gogama Forest Products and seasonal hunting and fishing by nearby residents, day visitors and individuals that base their activities out of local lodges and hunting/fishing camps.

The general area has been prospected for gold and base metals periodically since the 1910's. Numerous occurrences had underground development and production occurred at several deposits.

Four areas of the Platinex lands, as optioned from Skead Holdings, have been the subject of stripping, washing channel cutting and channel sampling. It is understood from discussions with the MNM and MNR personnel that no permits are required for the exploration work carried out to date with the exception of a 2 year renewable land use permit for a bridge installed by Platinex. Permits will be required from the Canada Department of Fisheries and Oceans and possibly the Ontario Ministry of Natural Resources if a disturbed area exceeds 2,500 square metres, drilling is done from lakes or if vehicles or equipment have to be taken across streams, creeks or rivers.

No attempt to evaluate any or all of the property for potential issues of environmental concern that may have resulted from either historic activities or the recent Platinex activities has taken place.

There is currently only one modern building on the subject property and the author is unaware of any associated environmental liabilities. Several remnants of old buildings were observed by D. Jamieson near the Churchill and Caswell shafts. Good fencing surrounds the Churchill shaft. The Herrick shaft location was expressed by a large hole that was essentially obstructed about two metres below the surface. The perimeter was marked by flagging since a barrier fence had not been present. The Caswell shaft was properly covered over and inaccessible.

4.0 COMMUNITIES AND FIRST NATIONS

4.1 Communities

The Platinex property is located between the regional centres of Timmins (192 km trucking distance) and Sudbury (206 km trucking distance). Both of these communities have extensive mining infrastructure.

The closest advanced gold mining projects to the Platinex Shining Tree property are the AuRico Young-Davidson project, located in Matachewan, 116 km trucking distance, the Lakeshore Gold Timmins West project, 172 km trucking distance from the centre of the Platinex property and the Trelawney (IAMGOLD) Cote Lake deposit, approximately 50 km trucking distance.

4.2 First Nations

The two nearest First Nation communities are located in the Matachewan area and Mattagami Lake, 20 km northeast of Gogama. An MOU has been entered into with the Mattagami First Nation. At this point in time, the Matachewan First Nation is involved in the process through the Mattagami First Nation.

Mattagami First Nation is an Oji-Cree community with a population of 450 members, 175 of which live on the Reserve. Several members of the community have prospecting and mineral exploration experience in the Shining Tree area. The band has an Economic Development and Employment Resource Centre on site.

The Matachewan First Nation is also an Oji-Cree community, with a population of approximately 60. In March of 2006, Northgate Minerals (now AuRico) and Matachewan First Nation signed a memorandum of understanding regarding development of the Young-Davidson gold project in Matachewan.

5.0 GEOLOGY

5.1 Quaternary Geology

The Quaternary Geology of the area was described by Roed and Hallett, 1979. An estimated 85% of the property is underlain by a ground moraine till with bedrock knobs and some recent peat and muck organic terrain. The area has moderate relief, is locally knobby and hummocky, and is generally dry. Local exceptions were noted in the extreme north of the main block and in the eastern half of the east block.

In contrast, the remaining 15% of the property at the eastern extremity is part of extensive glaciofluvial and glaciolacustrine deposits described as sand and gravel outwash plain as an apron about an esker with associated kame moraine ice contact stratified drift. These deposits are situated in an area of low relief and dry surface conditions. The extreme northeastern part of this claim block is similar with super-imposed recent peat organic terrain and a mixed wet and dry drainage.

Roed and Hallett (1979) reported that the Keewatin lobe of the Laurentide ice sheet advanced through the area at the beginning of the Wisconsinan (100,000 ybp) and that deglaciation of the area was completed by approximately 9,000 years ago. The authors classified the area as bedrock terrain with either exposed bedrock or bedrock covered by only 1-2 metres of ground moraine, whereas it may exceed 7 m thickness elsewhere. Glacial striae confirm the glacial advance was from the north.

The closest linear glacial features are north-south trending eskers located 8 km west and about 10 km east of the property. A section of the Sultan Scarp was reported about 3 km south of the property and was interpreted to be terminal moraine associated with a halt in the last Wisconsinan glaciation.

5.2 Regional Geology and Bedrock Assemblages

The Shining Tree greenstone belt is located in northeastern Ontario approximately 100 km north of Sudbury, and forms the southern portion of the Abitibi sub-province of the Superior Province (Figure 3). The Abitibi sub-province is one of the most prolific areas for gold and base metal deposits in the world. A strong spatial relationship exists between gold deposits and major regional faults. A number of world-class gold deposits occur in the Timmins camp and eastwards into Quebec, with all being associated with the Destor Porcupine Fault (Figure 4). Another principal trend of world-class gold deposits occurs along the regional structure identified as the Larder Lake–Cadillac fault (LLCF). Quite significantly the LLCF can be interpreted to extend through the Shining Tree area (Figure 4).

The metamorphic grade throughout most of the Shining Tree area is mid- to low greenschist facies (Oliver et al. 1999a, 1999b). Amygdules in volcanic rocks are filled with chlorite, carbonate or quartz and there is extensive saussuritization of feldspars in mafic flows and intrusions. The youngest sequence, the Timiskaming is less metamorphosed than the older assemblages. Higher grade, amphibolites facies contact metamorphism has been identified adjacent to the Miramichi and Togo batholiths (Ayer, 2000) (Figure 5).

5.2.1 Assemblages

The Pacaud, Deloro, Kidd-Munro and Tisdale assemblages (Figure 5) are dominated by volcanic supracrustal rocks which were formed before the first phase of deformation (Oliver et al, 1999b). Felsic volcanic units close to the presumed tops of the assemblages in the Shining Tree area have been dated by U/Pb techniques (Johns and Amelin 1999). The ages of the older three assemblages (Pacaud, Deloro and Kidd-Munro) indicate that the greenstone belt youngs to the northeast (Ayer, 2000).

The Pacaud assemblage is mainly composed of massive and pillowed basalts and is associated with minor spinifex or cumulate textured komatiites. The Deloro assemblage is dominated by felsic volcanic rocks and is capped in many places by chemical sediments, seen as banded chert and jasper. The Kidd-Munro assemblage is a varied assemblage dominated by tholeiitic basalts and komatiites, with minor felsic volcanic rocks, and the Tisdale assemblage comprises mafic flows and intermediate to felsic pyroclastics and/or volcanoclastics (Johns, 1999a).

The next youngest rocks are Keewatin intrusive rocks identified as the Miramichi and Togo batholiths which surround the Shining Tree area.

The next younger Archean assemblage in the area is the Timiskaming age volcanic and sedimentary rocks. Clasts of the older intrusive rocks are common within the Timiskaming assemblage. The Timiskaming assemblage has been divided into two lithostratigraphically distinct groups, the Indian Lake group (ILG) in the south and east, and the Natal group (NG) in the north and the west (Johns and Amelin 1999; Johns 1999a). The Natal group comprises volcanic pyroclastic rocks and flows usually of alkalic-type, and associated volcanoclastic sediments.

Unconformities seen between the Natal group and the underlying Keewatin sequences (Johns 1996, 1997, 1999a; Johns and Amelin 1999) are evidence of a hiatus in volcanic activity. The predominantly volcanic facies of the Natal group is found in southeast Natal Township and grades north and west into reworked pyroclastics, wackes and siltstones in Kelvin Township. The volcanoclastic rocks and associated pyroclastics are mainly tuffs, lapilli tuffs and tuff breccias. Timiskaming sediments are also located in a wide strip trending west across the northern portion of Connaught and Churchill Townships (Carter 1987; Johns 1996, 1997 and 1999a; Johns and Amelin 1999). The ILG sediments are described as texturally immature quartz-feldspathic arenites and wackes with minor conglomerates and rare mudstones. Detrital zircons were U/Pb dated as 2702 Ma.

Recent drilling has intersected an altered syenite intrusive on the property in the McBride area-hole MP12-01. The syenite is believed part of a syn or post-Timiskaming Group Archean intrusive alkalic pluton possibly linked genetically to the trachyte porphyries.

More detailed descriptions of the general rock types of the Shining Tree area were provided by Ayer (2,000) and are provided in the following text.

Komatiites and Associated Tholeiitic Rocks

In the Pacaud assemblage, komatiitic flows are confined to a small area close to the top of the sequence. In the northeast part of the Shining Tree area, komatiite flows interbedded with tholeiitic basalts cover an extensive area towards the top of the Kidd-Munro assemblage. The komatiitic flows demonstrate macro- and micro-spinifex textures, polysuturing and brecciation. Thin dunitic cumulate horizons also occur in flows at several horizons in the Kidd-Munro assemblage. These komatiitic flows tend to be black and talcose unless silicified or otherwise altered, in which case they weather to a pale grey colour (Johns 1996, 1997; Johns and Amelin 1999). The komatiites are commonly carbonatized and in a few locations in the Pacaud assemblage they contain abundant fuchsite, imparting a strong green colour to the rock. This alteration is also associated with silicification. The High Field Strength Elements, rare earth elements (REEs) and some of the major element oxides such as Al_2O_3 and TiO_2 are relatively immobile and appear to be unaffected by alteration.

Komatiites

The komatiites from the Pacaud assemblage all have high degrees of carbonatization, which can be identified on a microscopic level, some have a schistose texture and none have their primary texture preserved. All of the komatiites from the Pacaud assemblage are petrographically different from the younger Kidd-Munro (KM) komatiites. A significant difference can also be seen in the geochemistry between the komatiites of the Pacaud and the Kidd-Munro assemblages. KM komatiites are very similar to the aluminum-undepleted Tisdale-type komatiites in the Abitibi greenstone belt (Xie and Kerrich 1994).

Tholeiitic Volcanic Flows

Basalts and basaltic andesite in the Shining Tree area occur largely as pillowed or massive flows, but also include variolitic and spherulitic flows, feldspar phyric flows and amygdaloidal flows. The pillows are commonly well preserved and are of the order of 1 to 2 m long and 0.5 m thick, with fine grained margins, and a minor inter-pillow hyaloclastite. Alteration includes late carbonate veins and silicification. The tholeiitic basalts have been examined in terms of the assemblages within which they are located, but there is little geochemical difference between them. All of the basaltic units have unfractionated primitive-mantle normalized multi-element distributions.

Mafic to Ultramafic Sill

A mafic to ultramafic sill identifiable in the airborne magnetic survey maps as a long, linear feature spans Fawcett Township. It appears to be concordant with the volcanic rocks of the Pacaud

assemblage. The sill is terminated by the Granite Lake Fault near the western margin of Fawcett Township. This sill has been interpreted as komatiitic flows (Carter 1977) and more recently as a thin sill due to its placement within felsic flows (Johns and Amelin 1999). The rocks from this structure are pyroxene-phyric, coarse grained and have a range in MgO from 10 to 19 wt %. Ultramafic samples from the sill are similar to aluminum-depleted (with respect to average Neoproterozoic komatiites) komatiites from Boston Township in the Abitibi greenstone belt, displaying very low Al/Ti ratios, possibly due to elevated TiO₂ concentrations (Tomlinson et al, 1996c). The structure is the only example of light rare earth element-enriched and heavy rare earth element depleted ultramafic rocks in the greenstone belt.

5.3 Structural Geology

There are two main phases of deformation and associated metamorphism in the Shining Tree area (Oliver et al. 1999a, 1999b) with rocks older than 2.7 Ga having undergone two periods of deformation. There are multiple deformation zones in the older volcanic rocks in which gold has been found, especially in Macmurchy and Tyrrell Townships (Johns 1996, 1997 and 1999a). The younger, less metamorphosed, Timiskaming assemblage has undergone only one period of deformation (Oliver et al. 1999a, 1999b).

Major displacement of lithological units occurs across several north-northwest trending faults. Locally, the most prominent is the Michiwakenda Lake Fault that extends from south of Michiwakenda Lake and across the north part of the main block. The Herrick deposit is about 400 m west of and parallel to the interpreted fault (Figures 6 and 7). Other major faults located in the surrounding area, are the Jess Lake and Foley Lake Faults in Macmurchy Township, the Spider Lake Fault in south-western Tyrrell Township and the Elephant Head Lake Fault in Connaught Township. No gold deposits are known to occur directly within the fault zones, but there may be a clustering of gold occurrences in proximity to the Michiwakenda Lake Fault.

On a regional scale the Michiwakenda Lake Fault merges with the Mattagami River Fault (Figure 4). This fault system is responsible for a six kilometre sinistral throw of correlative rock units in the Shining Tree area and the South Porcupine Destor Break near Timmins. The west Timmins Mine of Lake Shore Gold is marginal to and west of the Mattagami River Fault. In the Shining Tree area the Herrick deposit is marginal to the fault and the LLCF is interpreted to trend through the Shining Tree property.

The importance of such regional faults to spatial association with major mineral deposits is demonstrated by the Montreal River Fault 'MLF' (Figure 4). It forms the confining association in the Cobalt Silver camp in the south is proximal to the Young Davidson Mine where the MRF crosses the LLCF, is proximal to the Pamour Gold Mine where the MRF crosses the South Porcupine Destor Break and in Kidd Township is proximal to the Kidd Creek VMS deposit.

Could the Michiwakenda Lake Fault be of equal significance?

There is also strong potential that northwest to west-northwest trending structures could provide an important control feature for gold occurrences. The apparent alignment of the gold occurrences from the Caswell deposit to the Kingston deposit in southwest Macmurchy Township can be projected toward the Herrick deposit which has north-south striking quartz-carbonate veins. There could be a complementary relationship between the various faults. Carter suggested a relationship between the gold-bearing structures and N50°W fold axes in south-western Macmurchy Township. Thus the Saville and Evelyn structures would be longitudinal faults, the N50E vein swarm on the Caswell would be cross-fracture fissure veins, and the Herrick, Ronda and Foisey structures would be diagonal shear veins.

A broad deformation zone was identified by Johns (1996) entering the property in the northeast and trending at roughly 065-085°. Displacements of the NNW trending magnetic features such as diabase along this deformation zone were noted by Dvorak (2008) as a regional structure. Sericite and chlorite schist within the volcanic units on the property along Highway 560 are evident. This is believed to be a shear zone which is either an extension of or a related splay to the LLCF.

The most significant known gold-bearing features in the Shining Tree area can be summarized as follows:

1. The Herrick, Ronda, Foisey, and Gold Corona quartz-carbonate veins and shear zones are oriented north-south.
2. The Evelyn, Saville (Caswell area), Gosselin and Discovery (Speed Lake area) quartz-carbonate veins and shear zone are oriented NW-SE. The Evelyn and Saville zones where intersected in 2011 drilling appear to the interflow sedimentary rocks whereas the Saville structure is recorded as a shear hosted quartz vein in historic records.
3. The Churchill, Pet, Speed Lake stripping (dominant parallel vein set) and the vein swarm at Caswell are NE-SW trending structures.

The most significant gold deposits known thus far in the Churchill-Macmurchy area of Shining Tree occur in the Herrick and Ronda vein systems, both oriented approximately due north. Preliminary structural interpretation for the sense of movement along vein/shears at the Caswell stripped area indicates that the maximum shortening direction is NNE-SSW, roughly sub parallel to the average orientation of Herrick and Ronda structures (Figure 6). Further examination of the structural components of all known showings and compilation of the structural measurements taken throughout the Shining Tree greenstone belt could produce exploration guidelines similar to those used for large wrench fault systems.

The results of the gold in till studies done on a property wide basis did not produce strong values in the vicinity and down ice from known deposits. Rather the gold in till anomalies appear to evidence previously unknown and probably more significant but unexplored gold mineralization.

6.0 MINERAL DEPOSIT TARGETS

Archean-age lode gold (greenstone-hosted quartz-carbonate veins) deposits and volcanogenic massive sulphide deposits are valid exploration targets on the Platinex Inc. Shining Tree property.

6.1 Gold Deposit Model

Greenstone-hosted quartz-carbonate vein deposits or mesothermal orogenic type gold deposits have been defined and described by Dube and Gosselin as follows: *“They correspond to structurally controlled complex epigenetic deposits hosted in deformed metamorphosed terrains. They consist of simple to complex networks of gold-bearing, laminated quartz-carbonate fault-fill veins in moderately to steeply dipping compressional brittle-ductile shear zones and faults with locally associated shallow-dipping extensional veins and hydrothermal breccias. They are hosted by greenschist facies to locally amphibolite facies metamorphic rocks of dominantly mafic composition and formed at intermediate depth in the crust (5-10 km). They are typically associated with iron-carbonate alteration. The mineralization is syn- to late deformation and typically post-peak greenschist facies or syn-peak amphibolites facies metamorphism. They are genetically associated with a low salinity CO₂-H₂O-rich hydrothermal fluid thought to also contain CH₄, Na, K and S. Gold is largely confined to the quartz-carbonate vein network, but may also be present in significant amounts within iron-rich sulphidized wallrock selvages or silicified and arsenopyrite-rich replacement zones. They are distributed along major compressional to transtensional crustal-scale faults in deformed greenstone terrains of all ages, but are more abundant and significant, in terms of total gold content, in Archean terrains....The*

diagnostic features of greenstone-hosted quartz-carbonate vein type gold deposits are arrays and networks of fault and shear-zone-related quartz carbonate laminated fault-fill and extensional veins in associated carbonatized metamorphosed greenstone rocks. The deposits are typically associated with large-scale (crustal) compressional faults, have a very significant vertical extent (≤ 2 km), with a very limited metallic zonation."

Dube and Gosselin also propose exploration parameters for quartz-carbonate greenstone-hosted gold deposits. They stated: *"In terms of exploration at the geological province or terrain scale, geological parameters that are common in highly fertile volcano-sedimentary belts include: 1) reactivated crustal-scale faults that focused porphyry-lamprophyre dyke swarms. 2) complex regional scale geometry of mixed lithostratigraphic packages. 3) evidence for multiple mineralization or remobilization events (Groves et al. 2003). The overprinting or remobilization was clearly a key factor in the formation of the exceptionally rich Goldcorp High-Grade Zone of the Campbell-Red Lake deposit (Dube et al. 2002; in press). The empirical spatial and genetic (?) relationship between large gold deposits and Timiskaming-like regional unconformity represents a first order exploration target as illustrated by districts such as Timmins, Kirkland Lake, and Red Lake (Hodgson, 1993; Robert, 2000; Dube et al, 2000, 2003 and in press)."*

6.1.1 Gold Potential of Platinex Shining Tree Property

The position of the Platinex Shining Tree property is shown by the red triangle on a depiction of the regional geology map by Dube and Gosselin (Figure 4). The relationship to major gold deposits, regional structures and the Timiskaming age sedimentary rocks are clearly favourable for gold exploration in this part of the Abitibi greenstone belt. Large gold deposits including world-class gold deposits have been found along regional trends which are shown in Figure 4 to pass through the Platinex Shining Tree property.

The Shining Tree property is located mid-way between the Young-Davidson and Cote Lake gold deposits along the interpreted western extension of the LLCF.

Gold occurrences on the Platinex Shining Tree property are spatially and likely structurally related, but have been worked historically as separate areas. Gold mineralization occurs in a variety of rock assemblages, rock types and structural styles in the Shining Tree area. The strong spatial association of the Ronda, Herrick, Churchill, Gold Corona, Buckingham and associated smaller gold occurrences implies a strong genetic association (Figure 6).

The Herrick vein system is controlled by what appears to be a cross-cutting structure that displaces Timiskaming age sedimentary and volcanic rocks. Alkali-rich porphyritic rocks (trachytic flows or intrusive) have been mapped and intersected in drill core at both the Herrick and Churchill prospects.

The Churchill vein system is located toward the top of the Deloro assemblage close to the unconformable contact with the Kidd-Munro assemblage to the north (Ayers 2002). Felsic fragmentals, cherty pyrite-rich sedimentary rocks, and sulphide facies iron formation are all associated with this assemblage contact. Work by Unocal in 1989 included geological mapping and diamond drilling of both the Churchill and Herrick vein systems. Timiskaming-aged volcanic rock types described by the field names "altered trachyte" and "red porphyry" were noted by Unocal in both the Churchill and Herrick vein systems and show a strong association with elevated gold values.

The Speed Lake or the Gosselin trend, which appears to be continuous from southeast Churchill Township into central Asquith Township, is located in the Pacaud assemblage. The Gosselin trend is closely aligned with a trend of ultramafic rocks, presumed to be composed mainly of komatiitic flows which have undergone deformation and alteration to form locally gold-bearing quartz-stockworks in fuchsitic carbonate alteration zones (green quartz-carbonate).

The Caswell vein system is part of a larger shear system that apparently trends southeast, similar to the gold-bearing Tyrrell shear zone further east near Gowganda. The host rocks are intermediate to mafic intrusive and extrusive volcanics which have undergone complex brittle to ductile deformation, are variably carbonate-altered and locally silicified with associated gold-bearing, locally pyritic, quartz-carbonate veins both parallel and discordant to the main shear orientation. In the author's opinion the N60°E to N85°E shear system at Caswell will prove to be a more important locus for gold mineralization. In contrast to other locations proximal to the historic large gold production areas, the Shining Tree area has received no comprehensive exploration campaigns and very little exploration below a depth of 250 metres.

6.2 Volcanogenic Massive Sulphide Deposits (VMS) Model

The secondary target type is volcanogenic massive sulphide deposits most commonly associated with the upper part of a mafic-felsic volcanic sequence of calc-alkaline affinity. Major deposits of this type have been mined in the Abitibi province, particularly in the Timmins and Noranda mining camps.

There is varying terminology for these essentially Cu-Zn-Pb-Ag dominated systems, such as volcanic-associated, volcanic-hosted, and volcano-sedimentary-hosted massive sulphide deposits. These deposits occur in submarine environments at or near the seafloor, as lenses of polymetallic massive sulphide. The mechanism for formation varies, but the key process is the discharge of hot, metal-bearing, hydrothermal fluids that have developed from sub-seafloor fluid convection.

VMS deposits are regarded as belonging to the larger group of exhalative deposits, which include iron formation and SEDEX deposits. Most VMS deposits consist of a tabular, stratabound accumulation of sulphides, quartz and accessory oxide and silicate minerals, stratigraphically above a discordant system of sulphide veins, and disseminations. Both the stratabound and discordant "pipes" have characteristic alteration halos which can extend significant distances from the sulphide accumulations, thus providing an important consideration in designing exploration programs for these types of deposits.

Another component of a VMS system is the heat source that provides the energy to develop the convection of fluids needed to scavenge and eventually concentrate metals. This generally takes the form of a subvolcanic intrusion, occurring some kilometres below the seawater interface. These heated intrusions are areally extensive, explaining the development of base metal "camps" consisting of numerous VMS deposits of varying sizes.

The process of delivering heated hydrothermal fluids into a column of seawater creates thin but regionally extensive iron-rich chemical sediment or "exhalites" that cap or are part of the VMS sequence. These units provide critical markers in VMS exploration programs.

6.2.1 VMS Potential of Platinex Shining Tree Property

The potential for base metal deposits in this area is noted by Unocal. Drill hole and surface observations in Unocal's Churchill area work, include strong sulphide mineralization in cherty horizons and pyritic, silicified argillaceous sediments, traces of sphalerite associated with pyrite, chloritization and weakly anomalous gold values (in 11.75 metres of "Sulphidic Rhyolite" in hole CU89-17). There are also anomalous widespread gold values within chert horizons (logged as 10.6 metres and 4.5 metres of chemical sedimentary rock, chert, sulphide iron formation in holes CU89-17 and 19). There is no copper, zinc, lead or silver analyses presented for any of the Unocal work. Whole rock and trace element data for eleven samples are presented, but Cu, Zn, Pb or Ag were not included in this data.

Regionally, these felsic and chemical sediment rocks intersected in the Churchill area occur at the top of the felsic volcanic dominated, 2730-2725 Ma Deloro assemblage, and at the base of the 2717-2712 Ma Kidd-Munro assemblage. Portions of this assemblage contact occur on the Platinex property between

the Churchill/Macmurchy Township boundary and an area just north of Perkins Lake. Major VMS deposits in the Southern Abitibi Sub-province are hosted by rocks with the following ages: Noranda – 2700Ma; Kam Kotia – 2707 to 2705Ma; Kidd Creek and Potter – 2717 to 2714 Ma. (Fyon et al 1992).

7.0 DEPOSITS AND OCCURRENCES OF THE SHINING TREE AREA PROPERTY AND LOCAL AREA GEOLOGY

7.1 Past Producers

There were no past producers on the Shining Tree property. Details about one nearby producer and two producing mines from the general area are provided in the following paragraphs.

7.1.1 Ronda Mine

The Ronda Mine, associated with the Ribble quartz vein is located in Macmurchy Township near Ribble Lake, a short distance southeast of the north part of the main claim block (Figure 6). The vein was reported to have a north strike and 60° west dip, was exposed almost continuously for 760 metres and had an average width of 1.5 metres (MNR, OGS Mineral Deposits Circular 18, Part 2, p. 75-76, 1979). The vein was described as intensely crumpled and folded and enclosed in carbonatized and pyritized schistose pillow lava. Intermittent development work occurred from 1912 to commencement of production in 1939. Production occurred only in 1939 and amounted to 2,727 oz. gold and 4830 oz. silver from 24,592 tons for an average grade of 0.11 oz. Au per ton in 1939. Plans of the workings indicate a westerly plunge to the mineralization which is projected to underlie the Platinox Shining Tree property at a depth of 300m.

7.1.2 Tyranite Mine

Further northeast of Macmurchy Township is the Tyranite Mine site, located a short distance south of the east-west boundary between Tyrell and Knight Townships. Graham (1932, p. 49, 51-52), The Northern Miner (1939) and Tremblay (1947) provided details about the setting and production. Discovered in 1930 as the Hedlund property, work by several companies culminated in 1939-1942 production of 31,352 oz. gold and 4,860 oz. silver from 223,810 tons of ore. The ore occurred as pods and lenses in two parallel, north-trending carbonatized and pyritized shear zones. Attitude of the shear zones was N20°W with a 70° west dip. The south ore body was 120 m by 12 metres and the north orebody was 60 metres by 8 metres wide. The shaft extended to 350 metres depth and development occurred on seven levels.

7.1.3 Bilmac Mine

The Bilmac prospect had production associated with the Evelyn and Saville Veins in 1922 and 1933. Total production has been reported as \$1607 for gold and silver combined (MNR, OGS Mineral Deposits Circular 18, Part 2, pp74-75, 1979).

7.2 Occurrences with Past Underground Development

The Herrick gold deposit and the Churchill and Caswell gold occurrences, located on the Platinox Shining Tree property, were previously investigated by underground workings. Elsewhere in the Shining Tree area there were nine additional occurrences that were also evaluated by means of underground workings. Details about the Herrick and Churchill deposits, and the Caswell area are provided in the following sections.

7.2.1 Herrick Deposit

The following timeline is based on information from Unocal (Cluff, 1989) and MNM sources.

- 1918: J.A. Knox discovered the Kingsley Vein
- 1918: Herrick Gold Mines acquired property; 4 holes drilled (955 metres), trenching and sampling; two compartment shaft to 15 metres.
- 1919: Report by F.V. Marsden M.E.; few details but recognized high gold values are wide-spread.
- 1920: Report by G. R. Rodgers.
- 1921: Report by P. McDonald for directors of Herrick Gold Mines. Noted that shaft had been sunk to 120' with limited lateral work; stations in shaft at 50 and 100' with 40' of drifting; also noted that surface sampling was done by cut channels every 5' was carefully done; discussion of free-milling metallurgy of ore; indicated extraction of over 90% of gold with fine grinding and cyanidation; low cyanide consumption.
- 1923: Work ceased with shaft at 300' and levels at 50', 95' (250' drifting), 200', and 300' (290' of crosscutting and 600' drifting).
- 1923: Henderson Bros. Mine contractors (F.D. Henderson) wrote a letter to F.C. Sutherland, King Edward Hotel regarding shutdown of Herrick operations. Felt the engineer of Tonapah Mining did not give the operation a chance and that the shaft should have been sunk to 500'. Believed sampling results were understated.
- 1933: Plans were made by Consolidated Ontario Gold Mines Ltd. to deepen the shaft to 700' in order to explore intersection of 0.16 opt over 50 feet. Plans were not carried out.
- 1933: Report by Kindle (Consulting geologist, Toronto). He noted 259' crosscutting and 631' of drifting on 300' level and that the vein was in the face of the last round according to last shift of workers. Drift at 100' level followed vein for 250'; ore found at 25' level; vein at shallow depth was offset to the west by a low angle fault passing through the shaft; Kindle took 11 samples, averaging \$10.20 /ton gold, similar to Marsdens report (average carried by a \$77.20 grab and \$13.40 across 2'; mainly grab sampling; recommendation for further underground development.
- 1935: Acquisition by Grantland Gold; compilation of data; Erie Canadian Mines drew 300' level plan dated April 15, 1935.
- 1935: G.D. Holbrooke report of April 15, 1935. A stamp indicates this report was received from Sylvanite Exploration Department in 1961 by the Resident Geologist for the ODM. The date, April 15, 1935 is also on two Erie Canadian Mines drawings (300' level plan and a composite section of four drill holes (original by G. R. Rodgers). The 300' level has approximately 900' of drifting and cross-cutting according to Erie Canadian Mines, 300' level plan April 15, 1935.
- 1940: Examination by Sylvanite Gold Mines; re-sampling of surface; no documentation of this located to date.
- 1962: Matachewan Canadian Gold Ltd., acquired property.
- 1969: Triton Exploration Ltd. acquired property.
- 1988: 751160 Ontario acquired property by staking.
- 1988-89: Unocal Canada Ltd. exploration program initiated, as they felt previous work indicated that the Herrick had the potential to host mineable reserves of 1732 tonnes per vertical foot at a

grade of 7.2 g/t Au over a 1.8m width with 50% dilution. Power-stripping, mapping, geophysics, channel sampling and diamond drilling (1473m in 11 holes).

1990: Fort Knox Resources examined the property as part of regional exploration work.

7.2.2 Caswell Occurrence

The following timeline is based on information from assessment files, Ministry of Northern Development and Mines sources and a report from Practical Exploration and Development Corporation (Edgar, 2001) (Figure 8).

- 1916: J. Messer reported on sampling that yielded numerous assays greater than one ounce per ton. Assays from a 40 foot shaft sunk on the east shore (No. 2 shaft) yielded assays up to 19 ounces per ton over 12 inches. No plans or maps for this data are known to exist.
- 1923: R.W. Demorest in 1923 reviewed work done on the property to date. Sampling by Demorest on 11 veins included assays from 0.12 ounces per ton over 30 inches to 31.16 ounces per ton over 48 inches.
- 1925: C. Baycroft reported on a diamond drill program by Canadian Champion Reef Mining Company Limited. A 92 foot deep shaft on the east side of the lake was also sunk around the same time, with 178 feet of drifting done from the bottom of the shaft.
- 1929: Canadian Champion Reef Mining Company Limited sank the No 1 shaft to 520 feet with 1105 feet of drifting; little documentation of sampling results available.
- 1933: Canadian Champion Reef Mining Company re-organized into Burvan Gold Mines Ltd.; re-sampling and prospecting by Burvan.
- 1938: F. Austin examined the data available for the Caswell property; noted lack of rigorous sampling correlation and documentation.
- 1975: New Bedford re-sampled several of the veins and drilled three holes. Visible gold was noted in two of the holes, but the best gold assay was 0.085 opt over 2.5 feet. Small bulk samples near shaft 2 (M. Newbury personal communication) returned gold assays of up to 2.96 ounces per ton and silver assays up to 1206 ounces per ton.
- 1980: Tut Explorations partially dewatered #1 shaft and performed limited underground sampling.
- 1987: Chesbar Resources examined the property and available data. 106 grab samples collected by Mike Perkins of Chesbar Resources. Eleven samples returned greater than 0.1 ounces per ton.
- 1988: Chesbar Resources drilled a total of 5,874 feet on veins 1 to 4 and 101 to 122.
- 2001: Practical Exploration and Development Corporation compiled available data on the property and performed limited power-stripping and sampling.

7.2.3 Churchill Occurrence

1918 – 1936: Exploration work resulted in a 38 foot deep pit on one of the veins and a 7 foot by 9 foot vertical two-compartment shaft to a depth of 110 feet on the north or No. 3 vein (Sinclair *et al.* 1935, p.82; Laird 1935, p.40). A level was set up at 109 feet on this shaft and 70 feet of drifting and 154 feet of crosscutting were done (Sinclair *et al.* 1936, p.92).

1989: Unocal Canada Ltd. acquired the property; performed geological mapping, power-stripping, channel sampling and drilled four holes for a total of 461 metres of diamond drilling.

7.2.4 Other Underground Development Sites

Two gold prospects with underground development, the Jefferson Lake and the Gosselin (Fred) prospects are located west of the Platinex Shining Tree property. Four other gold prospects, the Atlas, McIntyre-McDonald, Bennett and Kingston gold occurrences are located southeast of the Caswell shafts and were all evaluated with underground workings.

Most of the underground development work on these prospects dates from the early days of the Shining Tree gold camp. No major exploration programs have followed, but sporadic stripping, trenching, re-sampling and minor drilling have taken place during periods of high gold prices.

7.3 Occurrences and Prospects

7.3.1 On-Property

The McBride Royal Mining, Clarke and Knox gold occurrences have been documented on Ontario Geological Survey maps and in reports. All are shown on Figure 6.

Royal Mining Occurrence

Geological and geophysical mapping were carried out by Barringer Research Limited in 1971, outlining a shear zone in an east-west to east-northeast direction, immediately north of highway 560. (Carter, 1980).

Clarke Gold Occurrence

The Clarke gold occurrence is located in the extreme southwest corner of the property, in Asquith Township. Refer to Section 8.4 for details.

Knox Gold Showing

The Knox showing is located in Macmurchy Township, adjacent to Knox Lake in the east part of the east claim block. In the early days of the Shining Tree camp, this showing was known as the Wood claims. Hopkins described the occurrence as *“two narrow east-west rusty schist zones which contain a few parallel quartz veins, all being cut by a north-south diabase dyke. A little gold occurs in the north vein immediately east of the diabase and in the south vein directly west of the diabase.”*

Sampling done in 2004 and reported in the MDI for the Knox veins (two of 14 samples) returned assays of 0.029 and 0.018 opt from trench 1A and noted sampling by R.C. Whelan returned values up to 804 ppb.

7.3.2 Occurrences in the General Area

Other gold occurrences in the general area of the Platinex property include the Gunter, Gold Corona, Cochrane, Foisey, Featherstone and Onitap. Locations are shown in Figure 6. Details of these occurrences are beyond the scope of this report.

8.0 PROPERTY GEOLOGY

The Platinex Shining Tree property has undergone significant exploration in three separate areas: the Herrick Deposit, the Caswell area and the Churchill area. There is also a mineralized trend on the west boundary of the property, which has been designated as the Speed Lake area and more recent efforts have been made follow up highly anomalous gold in till results near Bielby Lake, Ribble Lake – McBride, the Clarke showing area and Cryderman Lake.

8.1 Herrick Deposit

The geology of the Herrick area is interpreted to be dominantly Timiskaming age. Regional mapping (Ayer 2000) indicated that a thick sequence of Timiskaming-age sediment is located just to the north of the area. Property scale work by Unocal Canada Limited revealed that alkalic extrusive/intrusive rocks (trachyandesites based on whole rock geochemistry) are intercalated with conglomerate and wackes and extend further south than previously mapped during regional mapping programs (Figures 6 and 7). The alkalic rocks are assumed to be related to the Timiskaming sediments further north. The stratigraphic relationship of a locally pyritic felsic fragmental unit located north of the main Herrick showing, in fault contact with greywacke, is unknown. The felsic fragmental is either part of the Timiskaming-aged sequence or is faulted or in-folded portion of the older Keewatin sequence. Strong silicification has been noted at the faulted felsic/greywacke contact, and was called a “silica cap” (Cluff, 1990). Another observation from drilling is that the Herrick veins appear to dilate towards the volcanic contact. Parallel or sub parallel to the sedimentary and trachyte units the sediment volcanic contact trends WNW and dips steeply south.

The most abundant rock types in the area of the Herrick deposit are clastic sediments, which occur either as wackes (greywacke) or conglomerate, with some gradation between the two units. Wackes have been described as either fine-grained, silty, and laminated with intervals of siltstone, and graphitic argillite, or as fine to medium-grained, massive, arkosic to pebbly. Greywacke can contain intervals of siltstone, argillite or graphitic sediment. Conglomerate is generally boulder or cobble clast supported, with a feldspathic gritty matrix. Clasts of granite, quartz, feldspar, felsic porphyry, felsic to mafic volcanic, siltstone, jasper, syenite, trachyte, silica cap breccias and pyrite have been noted within the conglomerate (Cluff, 1990 Figure 7).

Alkalic flows or sills have been noted in outcrop as more or less two continuous conformable southeast trending units between the conglomerate and/or wacke units. Diamond drilling by Unocal intersected several thick sections of “red porphyry”, as well as a few dyke or sill-like bodies. The mineralized Herrick shear/vein structure is often directly at a “red porphyry”-sediment contact. From surface mapping, it appears that this may simply be the effect of displacement of trachytic and sedimentary units along the Herrick structure creating structural contacts, rather than original contacts controlling mineralization. However, the use of the phrase “red porphyry” in the drill logs instead of trachytic volcanic rock indicates that drill core evidence observed by Cluff favoured an intrusive origin for the alkalic rocks. Although the trachyte porphyry is irregular in shape and provenience in many cases, it does tend to be parallel or sub parallel to the sedimentary units.

Small bodies of felsic porphyry were observed in drill core and were described by Cluff as grey feldspar porphyry or spotted feldspar porphyry. Grey feldspar porphyry is similar to the alkalic rocks, but with a carbonatized siliceous matrix, with spotted porphyry containing up to 1 cm diameter anhedral feldspar phenocrysts in a carbonatized siliceous matrix (Cluff, 1990).

Three southeast trending diabase dykes, approximately 200-300 metres apart, cut all rock types. The central diabase dyke cross-cuts the Herrick structure at a moderate angle (approximately 35 degrees), dips approximately 60 degrees southwest and is 10-15 metres thick. The diabase appears to truncate the north-south trending, gold-bearing shear zone that hosts the Central Zone. The parallel gold-

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

bearing, shear zone to the east hosts the Central Zone and appears to continue through and slightly offset the diabase dyke (Figure 7).

Sub-parallel to the diabase dykes are a set of late sub-vertical, faults that cause little horizontal displacement of stratigraphy/structure and an unknown amount of vertical displacement. One of these faults separates Timiskaming wacke from what appears to be Keewatin felsic fragmentals. The fault may also truncate or displace the Herrick northward extension of gold zones.

Gold mineralization at the Herrick deposit occurs along a north-south vein/shear structure. Movement along the structure is dextral and multi-staged, as evidenced by offset of an intruding diabase dyke along the Herrick structure. Timiskaming aged rock units are displaced up to 50 metres horizontally and an unknown distance vertically. Alteration consists of strong iron carbonatization, a moderate increase in potassium and a moderate decrease in sodium (Cluff, 1990). Sulphides are common in the 1-3% range, with minor arsenopyrite and tourmaline.

Quartz veins, stringers and breccias occupy the main 2 to 5 metre wide portion of the structure, as well as numerous narrow sub-parallel to strongly discordant fracture, shear and alteration zones. These smaller features locally contain gold mineralization. Larger splay features or parallel gold-bearing features have been designated as zones (Central Zone, West Zones and East Zones), such that the measure of the entire width of structure in which gold mineralization occurs is up to 80 metres, east to west (drill section 13+35N) (Figure 14). Platinex's drilling showed that the splay structures are attractive targets for follow up exploration achieving widths of 46m in hole HP10-44.

During the exploration by Platinex the state of the Unocal grid was examined. The Unocal grid was established in 1989 as a metric grid and all exploration done by Unocal and limited work by Fort Knox Gold Resources used this grid for reference. Very little of this grid is in evidence today around the power-stripped area. Some landmarks could be used to re-establish this grid, as a great deal of the important historic work is referenced to it. The grid is oriented north-south, with a 400m long baseline and cross lines at 50 metre intervals (total 4.83 line kms). However, there were some inconsistencies with the Herrick grid and hence it was decided to transpose and map all data on a UTM grid.

The claims immediately south of the original Unocal property are now part of the Platinex Inc. Shining Tree property. At the time of Unocal's exploration programs the claims were being explored by INCO. The INCO grid line terminations are shown on Unocal maps, and some of INCO's work has been filed as assessment reports. These reports show no indication of work done to physically trench or power-strip overburden to locate the south extension of the Herrick vein structure, despite the fact that Unocal stripping extended to within a few metres of the INCO claim boundary. There is a pencil notation on the Unocal geology map that some power-stripping has taken place just south of the Unocal/INCO boundary, but no results of this were observed or are known to be documented.

Two gold-mineralized veins were located by Inco along the edge of a 300 degree trending topographic low occupied by a creek and bay forming part of Michiwakenda Lake. One showing "consists of four separate 15 to 25 cm wide, bluish white, quartz veins hosted by massive intermediate volcanic rock. The veins are mineralized with 3 to 5% pyrite and chalcopyrite in up to 4 cm diameter blebs that are distributed sporadically throughout the veins. The veins occur in a parallel set spanning a width of 2 metres. The enclosed wallrock is moderately sheared parallel to the veins and is strongly chloritized. The vein set strikes at 260 degrees and dips 65 degrees to the southeast. A total of sixteen grab and chip-samples was collected from the veins and the chloritized wall rock" (Clark, 1989). No assays are provided in Clark's report. The second showing was located on the edge of a bay of Michiwakenda Lake. A one to two metre wide "grey white quartz vein outcrops on the shoreline....striking 300 degrees with a vertical dip; it is hosted by moderately sheared and chloritic intermediate volcanics. This vein is sporadically mineralized with 5-8% pyrite and chalcopyrite as isolated blebs to 2 cm in diameter. A total of fourteen grab and chip-samples were collected from the vein and wallrock material." No assays for

this showing are provided in the report (Clark, 1989). The two showings occur 500m south of the southern-most exposure of the Herrick vein structure.

One till sample collected south of the Herrick deposit (North 5273111; East 484084; UTM NAD83 Zone 17) some 200m returned a total count of 25 gold grains. It is speculated that the source is actually east of the Herrick Central zone. It is noteworthy that the Herrick Central zone exposing many tens of metres of strike length with high grade gold does not appear to have a highly anomalous gold in till signature.

8.2 Caswell Area

Geology of the Caswell area is dominated by Pacaud assemblage mafic volcanics and related subvolcanic diorite and gabbro. Quartz porphyrys were noted by early workers in the area, but were not observed by Platinex geologists. A distinctive feature of this area is the prolific development of quartz veins that occur on both sides of the West Shining Tree Creek often called Caswell Lake (Figure 8).

Veins on the east side of the creek have been numbered by previous workers from 101 to 124. Veins on the west side of the creek have been numbered from 1 to 17. No sampling plans have been located to define the extent and location of gold values within individual veins. There were numerous sampling efforts by mining companies, geologists and mining engineers between 1916 and 1936. Little of the data is properly documented with plans, sections, or assay certificates. To date the most comprehensive review of the property is given by Bruce Edgar in a report for Practical Exploration and Development Corporation in 2001 and submitted for assessment credit. The following summary is largely derived from Edgar's work, as well as MNDM annual reports and OGS open file reports.

The earliest available data was from J. Messer in 1916. He indicated that the No. 1 vein (renamed 7a vein) could be traced E-W for 1500 feet, averaged 1.0 feet in width and had 2.5 feet of sheared material on either side. Multi-ounce per ton gold grades were reported from sampling of this vein with widths up to five feet on both sides of the creek. Later workers did not assume the east and west veins were contiguous across the creek. Assays from a 40 foot shaft sunk on the east shore portion of the vein also returned very high assays up to 19 ounces per ton over 12 inches. The No. 2 vein was traced for 500 feet and returned gold values of 2.26 ounces per ton over 31 inches, 0.293 ounces per ton over 23 inches, and 0.767 ounces per ton over 78 inches. Sampling of the northwest trending Saville vein (also named the No 4 vein), where exposed near the south boundary and shoreline, returned an average of 1.49 ounces per ton over 33 inches. No plans or maps for this data are known to exist.

R.W. Demorest in 1923 reviewed work done on the property to date and described the Saville or No. 4 vein as being traceable for 1.5 miles at N 46°W and dipping 65°W. A parallel shear zone 20 feet wide located beneath West Shining Tree Creek may have been a possible extension of the northwest trending Evelyn Vein that was explored to the southeast on the Bilmac property. Sampling by Demorest on 11 veins included assays from 0.12 ounces per ton over 30 inches to 31.16 ounces per ton over 48 inches.

In 1925, C. Baycroft reported on a diamond drill program by Canadian Champion Reef Mining Company Limited that tested the Saville structure beneath Caswell Lake, as well as east-northeasterly striking veins on the east side of the lake. Six of the seven holes reported high grade gold values over 5 and 10 foot widths, and included 5.40 ounces per ton over 5 feet, 4.63 ounces per ton over 10 feet on the Saville structure, and 0.61 ounces per ton over 5 feet and 5.7 ounces per ton over 5 feet on veins on the east shore.

The 92 foot deep shaft on the east side of the lake was also sunk around the same time, with 178 feet of drifting done from the bottom of the shaft.

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

Canadian Champion Reef Mining Company Limited then carried out development work, beginning in 1928, on the west shore of the lake. A shaft was sunk to 520 feet (No. 1 shaft), with levels at 100, 250, 375 and 500 feet, with lateral development of 325 and 780 feet on the 250 and 500 foot levels. Unfortunately there is very little record of the sampling results from this work.

By 1933 Canadian Champion Reef was re-organized into Burvan Gold Mines Ltd. and work was done to de-water the No. 1 shaft and re-sample the workings, correlate surface and underground workings, and follow-up on the favourable drilling program conducted in 1925 (Laird 1935). Burvan also continued to prospect the surface veins, and located additional veins.

In 1938, F. Austin examined the data available for the Caswell property and questioned the reliability of both the drilling and underground work. He pointed to a lack of systematic underground sampling and mapping and lack of surface drilling to follow up the initial results obtained in 1925. Apparently only assays for work on the 100 foot level were available. Drifting on this level returned "good results" for 60 feet. One ton of ore treated by the Ontario Department of Mines from another section with quartz stringers and visible gold returned a gold value of 0.27 ounces per ton.

There was no additional work done on the property until 1975, when New Bedford Explorations re-sampled some of the trenches and drilled three holes. Visible gold was noted in two of the holes, but the best gold assay was 0.085 opt over 2.5 feet. Small bulk samples near the Number 2 shaft (Mike Newbury, personal communication) returned gold assays of up to 2.96 oz/t and silver assays up to 1206 oz/t.

In 1980, Tut Explorations de-watered and repaired the No. 1 shaft. The second level (250') was mapped and sampled. High grade samples were said to be obtained, but no assay plans are available.

In 1987 Chesbar Resources examined the property and available data. There is a mention that Falconbridge completed geophysical surveys and 2,434 feet of diamond drilling in four holes directed at the Saville structure. No record of this work is available. Mike Perkins of Chesbar Resources collected 106 grab samples and eleven samples returned greater than 0.1 oz/t Au.

A 5,874 foot drill program supervised by Perkins, tested veins No. 1, No. 2, No. 3 and No. 4 as well as most of the veins on the east shore (No. 101 to 122). Significant gold values were returned from the No. 3 Vein (0.152 oz/t over 6 feet) and the No. 101 Vein (0.098 oz/t over 5 feet).

8.3 Churchill Area

Southwest of the Herrick area are a number of showings which appear closely related to the contact between the Deloro and Kidd-Munro assemblages. These showings include the Cochrane, Gold-Corona and Churchill vein/alteration systems. The Churchill veins are covered by the Platinex Inc. Shining Tree property, as is part of the Deloro/Kidd-Munro contact (typically mapped as chemical sediments) further to the northwest.

Laird visited the property in 1933 and reported visible gold in many places along the 30 metres of exposure in the No 1 vein and visible gold at the west end of the No 3 vein. In a 1934 report he stated: *"The south vein, known as No. 1, and that part of No. 3 vein east of the fault occur in a light-coloured rhyolite or quartz porphyry, which is somewhat sheared at S80°E. This porphyry, which is probably a differentiated portion of the large granitic mass lying to the south, occurs as a narrow tongue pinching out a few hundred feet to the north of the showing. Locally, it is intersected by irregular stock works of narrow quartz stringers, which are reported to carry low values in gold. The veins are of the fissure type and are definitely later than the porphyry, since they cut it. The north-south faults displacing the veins belong to a late system of north-south fracturing characteristic of the whole region. In the case of the adjoining properties, (Gold Corona, Herrick, and Wasapika) the north-south fractures carry the gold-*

bearing quartz veins, but on the Churchill property the east west fractures are in this respect the important ones.”

Laird also compiled the company assay plan and indicated that channel sampling of the No. 3 vein over a 90 metre exposure returned an average gold value of 27.5 g/t over a width of 1.2 metres and that a 30.5 metre exposure of the No. 1 vein returned a gold value of 29.4 g/t over a 1.2 metre width.

Unocal acquired the property in 1989 and compiled historic information, performed ground magnetic and VLF EM surveys and power-stripped the No. 3 vein for 95 metres and the No. 4 vein for 25 metres for a total of 205 linear metres. Forty-eight channel samples and six composite chip/grab samples were collected and assayed, and four diamond drill holes (461 metres) were drilled to test the property.

The geology in the vicinity of the old workings was mapped in detail by Cluff for Unocal Canada Ltd. The dominant rock type was felsic fragmental rocks, with associated crystal tufts and cherty or graphitic, pyritic sediments and minor clastic sediments. Mafic to ultramafic rocks occur within the felsic package as dark green black massive units. Whole rock, petrography and microprobe data were used to verify rock types. Some of the mafic volcanics were interpreted as altered feldspar porphyry of dioritic composition from this work (Cluff, 1990).

Moderate to strong alteration and shearing of the felsic volcanic unit created zones of banded pink and yellow schists and mylonitic rocks. Clots of pyrite occur sporadically throughout the felsic fragmental rocks.

A chemical sediment horizon, that appeared to correlate with sulphide iron formation mapped on properties further to the northwest, is located in felsic fragmental rocks just to the north of the vein system and consists of alternating bands of chert, massive pyrite, black argillite and graphitic schist up to 10 metres wide. No base metal minerals were observed. The geology is interpreted as to be indicative of a significant exhalative event wherein surrounding felsic fragmental rocks were sericitized and chloritized.

8.4 Speed Lake area

The Speed Lake area is interpreted to be underlain by Keewatin aged rocks of the Pacaud assemblage, that is, calc-alkaline and tholeiitic volcanic rocks and minor komatiitic rocks dated between 2745-2730 (Figure 6). A southwest trending feature which includes an extensive quartz vein system, altered mafic-ultramafic rocks (iron-magnesium-carbonate-fuchsite) and altered felsic intrusive and extrusive rocks (quartz-sericite-chlorite-epidote) has been termed the Gosselin Vein or “Gosselin Rift Zone” (McCannell, 1975).

The “Gosselin Rift Zone” has been described as a zone of faulting, fracturing and shearing approximately 200 metres wide, trending 150 degrees and dipping sub-vertically. East-west to east-southeast cross structures offset the zone, apparently by as much as 400 metres just south of Speed Lake. Gold mineralization tends to be associated with an increase in pyrite mineralization from trace levels to 4-5% in carbonatized volcanic rocks. Felsic volcanic rocks tend to be altered to sericite-epidote-chlorite, while mafic to ultramafic volcanic are dolomitized, with fuchsite (green carbonate), or have been silicified.

The eastern margin of the “Gosselin Rift Zone” is exposed on the western boundary of the Platinex property in Churchill Township, just south of Speed Lake. This area was the focus of a power-stripping program by Platinex Inc. in 2008 (See section 10.2 Speed Lake on current work – Speed Lake).

Previous exploration work indicated that green carbonate rock, quartz veins and iron carbonate alteration occur along the southeast side of the Gosselin Zone, and locally trend onto the Platinex Shining Tree claims in Asquith Township. Further to the southeast, a zone of quartz- carbonate

alteration and gold mineralization extends along an ultramafic unit mapped by Johns (Johns 2000). This trend hosts the Thompson-Peterson and Clarke gold occurrences on and close to the Platinex property and hosts the Buckingham gold deposit at the southeast extent, approximately two kilometres south of the Platinex Inc. claims.

Drilling by Asquith Resources in 1989 on the Buckingham property intersected gold values in “a very distinctive pale green carbonate rock which had been intensely sheared and injected with grey, blue, and black quartz veins. The shear zones may be up to 20 feet wide but the central core normally contains the higher gold values. Disseminated pyrite was pervasive throughout the shears, and increased in quantity with intensity of shearing. Gold was noted as minute specks along fracture zones in quartz veins. Tourmaline was common in quartz.

Among the better intersections in parallel sheared green quartz-veined carbonate zones:

“C zone” 8.9 feet of 0.217 oz/t and 4.0 feet of 0.099 oz/t

“D zone” 8.9 feet of 0.507 oz/t and 8.5 feet of 0.217 oz/t

“E zone” 4.0 feet of 0.137 oz/t and 7.0 feet of 0.118 oz/t

Intersections of a grey white quartz vein (without an associated strong carbonate shear) included:

8.4 feet of 0.275 oz/t and 2.4 feet of 0.437 oz/t.

The Clarke occurrence is on the southwest corner of the Platinex property in Asquith Township and is described as iron-magnesium-calcium carbonate with pyrite and quartz stockworks, some of which contain free gold (Gorden et al, 1979). Hopkins (1920) described the occurrence as “rusty weathering green carbonate, impregnated in places with iron pyrites and containing a network of quartz veinlets. Gold could be seen in some of the quartz stringers. The rock is an iron-magnesium-lime carbonate, with considerable silica, aluminum and a trace of nickel...on the northern part of the claim the carbonate is cut by red and grey granite-porphry and near the contacts are large lenses of quartz somewhat similar to those on the Gosselin and carrying feldspar, galena, chalcopryite, pyrite, talc and sometimes gold and other minerals.” Johns (2003) mapped dunitic komatiites in contact with a sill like gabbro at this location.

The Thompson-Peterson occurrence is north-northwest of the Clarke occurrence, but is just west of the Platinex Inc. claims. The description is similar to the Clarke occurrence: a zone of rusty iron-magnesium-calcium carbonate rock near a granite porphyry body, cut by a network of quartz stringers containing visible gold.

9.0 MINERALIZATION

The dominant style of gold mineralization on the Platinex Shining Tree property is shear-hosted quartz-carbonate veins, generally associated with pervasive iron carbonate or green carbonate alteration and low levels of sulphidization.

9.1 Herrick

Gold mineralization at the Herrick Deposit is associated with quartz veins, pyrite, tourmaline and arsenopyrite, with better gold values often occurring in pyritic quartz vein breccias. The quartz veins occur along a north-south trending zone of high strain that has displaced trachytic volcanic rocks and clastic sedimentary rocks, and provided a conduit for hydrothermal fluids. Exploration work by Unocal Canada Ltd. delineated this high strain zone along a strike length of 300 metres and a maximum width of 80 metres.

Several types of porphyritic intrusive/extrusive rock types were delineated by Unocal. The main units of interest are described as trachytes in surface mapping or red porphyry in drill logs. The trachyte or red porphyry is assumed to be an extrusive volcanic or synvolcanic intrusive sill that has suffered metre to 10's of metre scale movements along earlier north (Herrick vein structure) and later northwest and west-northwest trending faults. The trachytic rocks are observed to have undergone high strain, strong carbonate alteration, strong to intense quartz veining, and localized mineralization of pyrite, tourmaline, green mica, and minor arsenopyrite.

9.2 Caswell area

Gold is hosted by narrow quartz veins and lenses along shears oriented in a variety of directions, but generally either in a northwest or east-northeast direction. These veins exhibit both compressional (drag folds) and tensional (boudinage) strain. Quartz veins generally contain minor to 2% pyrite and local dark seams of chlorite and possibly very fine-grained tourmaline. Wall rock to the veins is often sheared and fractured with strong iron carbonate alteration extending for several metres away from the vein. Visible gold has been noted in the past, but was not observed by Bryant and Jamieson (2008).

The Saville vein occurs as part of the regional northwest shear structure which hosts the Caswell veins and numerous occurrences to the southeast (Bilmac, Bennet, Atlas etc.). Where exposed on the Platinex Shining Tree property on the edge of Caswell Lake, the vein is up to a metre wide with quartz stringers developed in adjacent iron carbonate-rich schist. Anomalous gold values were returned from chip and channel sampling done on the vein by Platinex in 2008. Drilling by Platinex in 2010 tested the Saville and Evelyn veins severally revealing that both appear to be graphitic and pyritic interflow sedimentary units where tested (Figure 8).

9.3 Speed Lake area

Gold is hosted by quartz veining and stockworks within green carbonate or iron carbonate altered mafic to ultramafic volcanics. Sulphides are generally a minor component of green carbonate-hosted mineralization, but are common in veins within more mafic rocks. A few hundred metres west of the Platinex Shining Tree property, the Gosselin vein strikes 150 degrees and has been traced along strike for several kilometres. It consists of white quartz-carbonate material with sporadic sulphide and gold mineralization. In places the vein is greater than 10 metres wide (Carter, 1980).

10.0 EVALUATION OF SHOWINGS

Exploration activities during the summer of 2008 included detailed evaluation of showings at Caswell and Speed Lake. Stripping on Caswell was also done in 2010 and stripping south of Beilby Lake was done in 2011. Caswell was drill tested in 2011 and Beilby Lake was drill tested in 2012. Details are provided in the following sections.

10.1 Caswell area

In May and June of 2008, Platinex Inc. and Robert Macgregor conducted overburden stripping, washing and channel sampling on claim L4203531 in Macmurchy Township south of Michiwakenda Lake. The centre of the stripping is at 485940E and 5272055N (UTM Zone 17 NAD 83), with an exposed outcrop area totalling 1600 square metres.

The area stripped is a prominent hill southeast of the muck piles near the Caswell No. 1 shaft area. Previous trenching in the area had revealed a number of quartz veins from which high gold values had been obtained.

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

The 2008 stripping program exposed the historic No. 1, No. 2, No. 3 and No. 4 or Saville veins. The veins follow narrow (0.5 to 1.0 metre) fractures and shear zones, trending northeast or west southwest, within a locally carbonatized intermediate intrusive rock resembling diorite or leucogabbro. Moderate to strong carbonate alteration extends from the vein/shear structures for several metres into the wall rock. Pyrite is common in proximity to veining, but is not pervasive or abundant. The No. 4 or Saville vein is hosted by mafic volcanics and a volcanic fragmental unit exposed at the base of the hill on the shore of west Shining Tree Creek.

Eighty-seven channel samples, six chip samples and one grab sample were taken and analyzed for gold at ALS Chemex in Vancouver. The best assay results were from the No. 1 vein: 11.05 g/t over 0.75m, 10.45 g/t over 0.5m, 5.99 g/t over 0.7m, 5.91g/t over 0.6m, 4.22 g/t over 0.75m. One sample related to the No. 2 vein assayed 9.11 g/t over 0.5m. Twelve additional samples assayed greater than 200 ppb. Sieve metallic analysis of ten samples showed two samples with a significant contribution from the plus fraction, increasing final reported grade between 8 and 25 % from the minus fraction pulps.

In October and November, 2010 Platinex conducted overburden stripping, washing and channel sampling on claims in Macmurchy Township south of Michiwakenda Lake and east of Caswell Creek. Quartz veins #101a, 101, 102 near shaft 2, 108, 109, 109a, 109b, 110, 111a, 115, 123 and L122 were exposed in eight trenches covering a total area of approximately 3,000m (Figure 8). The work covered nine of thirty-nine veins identified in historic work. M. Spaho mapped the showings, Katrine Exploration washed and collected channel samples marked out by M. Spaho.

The veins follow narrow 0.5 to 1.0m fractures and shear zones, trending 065° and 085° within locally carbonatized and sericitized intermediate to mafic metavolcanic and diorite. Moderate to strong carbonate alteration extends from the vein/shear structures for several metres into the wall rock. Pyrite is common proximal to veining.

In total 370 channel samples, and 4 grab samples were taken and analysed for gold by ALS Chemex of Vancouver by fire assay.

Individual channel samples returned numerous highly anomalous gold values including 10.55 g/t Au over/ 0.69m, 9.1 g/t over 1.0m, 4.69 g/t over 0.69m, 4.1 g/t over 0.44m, 4.41 g/t over 0.47m, 3.37 over 1.03m, 3.57 over 0.54m in channel samples and up to 27.7g/t Au in grab samples. In addition to these samples, 22 channel samples returned gold values greater than 0.7 g/t, three of which were greater than 2 g/t. Anomalous silver and tellurium values were evident throughout the area sampled, with assays up to 2.15 g /t Ag and 1.1 g/t. The gold enriched areas appear to be marginal to and probably beneath areas of anomalous silver-tellurium values, indicating the potential for Ag and Te to be important pathfinder elements.

In March and April, 2011 Platinex completed a seven hole 1,070m drilling program on the Caswell area (Figure 8). The first hole, WP11-01, returned a gold assay of 18.75 g/t over 0.5m within a broader zone of visible gold bearing quartz-tourmaline veining assaying 4.52 g/t over 2.52 m believed to be within vein 108 and immediately south of trench 4.

The remaining six holes focused on other shear zones, generally in locations that could not be easily accessed for trenching due to proximity to West Shining Tree Creek. A total of eleven separate quartz tourmaline vein/shear structures with anomalous gold values between 0.2 and 1 g/t were intersected, with total zone widths ranging from 1.5 to 22.5 metres.

Six of the holes were surveyed by DGI Geoscience to provide structural orientation of veins and shears observed in the drill holes. This technique will help distinguish the numerous vein sets on the Caswell prospect. In addition, an I.P probe and natural gamma log probe were used to provide data on

chargeability, resistivity and alteration (dominantly potassic). A broad alteration system controlled by a nearly orthogonal conjugate fault system is evident. The current interpretation is that each of the zones outlined above is a separate structure within the overall conjugate fault pattern. The 2011 drilling along with the 2010 trenching programs have tested a small number of structures within an area of 700 m by 400 m.

These results helped explain some of the previously described highly variable but widespread distribution of the gold within the area comprising 39 documented shear-hosted veins within an overall broad (800m plus) east-north east (070°) trending structure. In addition the previous grab sampling by Chesbar Resources and Practical Mining can now be put in context and selected results are shown on the attached map (Figure 8)

Two till samples were collected on the Caswell prospect returning highly prospective gold counts of 69 grains and 19 grains. The sample taken down ice but near trench 1 is dominated by pristine grains indicating a probable nearby source possibly north of the known explored portion of Caswell veins.

The Caswell area has been noted for some spectacular gold results flanked by a low tenor of gold mineralization. It would appear that highly strained and altered areas hosting gold are limited in lateral extent and no continuous potentially commercial gold grades were noted. However, the pathfinder associations such as tellurium and silver are very anomalous in all of these veins and provide evidence of the systematic nature of the mineralization. (Figure 8). Casaceli (Personal Communication, 2011) stated that the Tyrante Mine has similar associations which he believed are common to gold-alkaline systems.

10.2 Speed Lake Area

In May and June of 2008, Platinex Inc. and Robert Macgregor conducted overburden stripping, washing and channel sampling on claim L1242934 in Churchill Township south of Speed Lake and north of Highway 560. The centre of the power-stripping is at 480618E and 5269978 (UTM Zone 17 NAD 83), with an exposed outcrop area totalling 1100 square metres.

Old caved trenches, provided evidence of much earlier work in the area. A number of exploration programs since the 1912 discovery of the gold-bearing Gosselin vein system, had examined the area just west of claim L1242934.

The 2008 power-stripping program exposed ultramafic rocks with strong green carbonate alteration and significant areas of strongly developed quartz veining and quartz stock works. Traces of sulphide mineralization were observed, along with minor areas of pervasive silicification. The northern most outcrop in the trench was relatively fresh mafic volcanic in low angle fault contact with the green carbonate altered ultramafics.

Fifty-eight channel samples and seven grab samples were taken and analyzed for gold at ALS Chemex in Vancouver. The best assay results were 4.58 g/t over 2.5 metres, 3.91 g/t over 1.2 metres and 1.74 g/t over 1.0 metres. Three other samples returned values over 200 ppb with no significant results from the remaining channel and grab samples. Spatial clustering of these results suggests that a rubble-filled fault zone may control mineralization. Although some screen metallic assaying was done on selected channel samples, no such assaying was done on the samples with higher gold values. As a check the rejects of samples assaying greater than 200 ppb should be sent to a second lab for screen metallic analysis to confirm gold mineralization and understand the size distribution of gold grains within the samples. Further removal of rubble from fault-controlled depressions would facilitate follow-up channel sampling.

10.3 Beilby Lake Area

Follow up prospecting and stripping up ice direction from two high gold in till sample results (71 total gold grains in sample 54787) south of Bielby Lake resulted in the location of an area of elevated gold values in outcrop (Figure 9). Grab sample values to 0.75g/t Au were obtained and later channel samples recovered values of up to 2.65 g/t Au/ 0.56m.

The area exposed is underlain by well preserved unstrained variolitic pillowed basalts with modest hyaloclastite, a narrow unit of felsic volcanoclastics both intruded by an altered moderately to strongly magnetic gabbro. A 070° 0.5 – 1.0m thick rusty carbonate shear zone cuts and appears to sinistrally displace the various rocks approximately 20 metres. A NNW shear in turn displaces the 070° shear dextrally approximately 10m. The best value in channel sampling which was keyed to the 070° shear was 2.65 g/t Au/0.56m. Due to the widespread anomalous gold values and the high gold in till values which had not yet been adequately explained it was decided to cover the area with pole-dipole spectral IP and drill higher priority anomalies.

Lines for the IP survey were run at 200m intervals in a north-south orientation tied to the Herrick grid with a tie line at 8+00S crossing on the south side of the Bielby Lake exposure very close to the till sample location. Line 8+00W is within 15m of the outcrop on the east and line 10+00 W is approximately 115m west of the outcrop.

Hole BP12-01 was collared at 10+40W on the line 8+00N drilled due west to test a strong chargeability anomaly with high resistivity close to surface and low resistivity at depth. The hole intersected variolitic pillowed mafic volcanics from top to bottom of the hole at 170m. The rock is mineralized with fine grained pyrite throughout chiefly in selvages. The pillows are apparently unstrained. Due to a budget shortfall detailed logging and sampling were deferred.

Hole BP12-02 was spotted at 9+00S on line 8+00 W and drilled due north to intersect a strong IP anomaly. The hole intersected a sequence of predominantly felsic volcanic flows and fragmentals contacted by mafic volcanics and ending in diabase and trachyte. The felsic volcanics are intensively sericitized and hematized with intense quartz carbonate flooding over an 11m zone. Sulphides are disseminated throughout with stronger mineralization associated with quartz carbonate veining including trace chalcopyrite. The rock is relatively unstrained, however, and the assay results were low.

11.0 GEOPHYSICAL SURVEYS

11.1 Regional Gravity Surveys

The Ontario Geological Survey supervised and edited the compilation of digital gravity data obtained from the National Gravity Data Base maintained by the Geophysical Data Centre, Geological Survey of Canada (Gupta, 1991). The reconnaissance-scale data indicated that most, if not all of the property correlates with the middle portion of a composite gravity anomaly. The highest gravity responses extend northwest from the community of Shining Tree and from the east part of Macmurchy Township into Fawcett Township.

Gravity anomalies indicate thicker sequences of earth's crust where there would be a greater potential for partial melting and development of convection cells that could cause leaching of gold from thick volcanic, sedimentary and plutonic sequences, migration of the gold along structural conduits and the subsequent deposition of the gold in areas such as the Shining Tree property.

11.2 Airborne Geophysical Surveys

11.2.1 Prior Surveys

The Ontario Geological Survey commissioned an airborne magnetic and time domain electromagnetic (TDEM) survey which was flown for the Ontario Geological Survey by Geoterrex in 1990. The survey was completed with a fixed wing configuration that included a magnetometer system and GEOTEM II TDEM system. Flight line separation was 200 metres and the regional survey area included the Platinex project area.

Features interpreted from the total field data are:

- A magnetic anomaly apparently coincident with the diabase dike at the Herrick deposit. The feature extends somewhat intermittently south through the area just west of the Ronda Mine, on to the south-southeast beyond the property.
- Stronger magnetic response typical of the southwest part of the property is associated with mafic to ultramafic volcanic rocks.
- A magnetic high and flanking magnetic low in the area northwest of the Herrick deposit appear to reflect a folded feature associated in large part with the Timiskaming sediments identified by Carter and by Johns.
- Magnetic lows along most of Michiwakenda Lake and on to the south-southeast suggest the presence of the major fault.
- Distinctive, narrow north-northwest trending magnetic highs that cross the east block of the property likely indicate unmapped diabase dikes and likely a formational contact between mafic to intermediate and intermediate to felsic volcanic rocks.
- The northwest trend of the showings that had underground development, including the Caswell deposit, correlate with a magnetic low.

The second vertical derivative maps provided improved resolution of the features described above and also indicates:

- Significantly improved delineation of the individual lithological units identified by Carter and generalized by Johns.
- Indications that northeast trending faults or shear zones may be common.
- A common, pattern of apparent discontinuity of magnetic features caused by the wide-spaced flight lines.

The electromagnetic data from the survey indicated two conductive features apparently associated with bedrock and one anthropogenic-related anomaly that coincides with the community of Shining Tree and Highway 560. The largest electromagnetic anomaly extends about two kilometres with only minor segments associated with the west-central part of the property. It appears to be correlated with Churchill, Gold Corona to Cochrane showing and deposits which appear to be associated with the top of the Deloro assemblage and the adjacent intermediate to felsic intrusive. The potential for sulphide related gold mineralization and for base metal volcanogenic deposits both need to be considered. The second airborne EM anomaly appears to be a single-line anomaly located adjacent to the shore of Michiwakenda Lake near the north end of the property.

The overall impression was that more detailed airborne magnetic data would provide significant improvement in resolution to facilitate more accurate interpretation of structural features as well as lithological units that may host or coincide with many of the gold-bearing zones.

11.2.2 Platinex Airborne Survey

In July of 2008, Platinex contracted Terraquest Inc. to conduct a fixed-wing airborne survey on the Shining Tree project. A total of 491 line-kilometres were flown with the following sensors:

- High resolution aeromagnetic
- Horizontal magnetic gradiometer-
- XDS VLF EM
- Radiometrics (gamma ray spectrometer)

The nominal line spacing for the 050/230 degree oriented traverses was 100 metres, with nominal aircraft clearance of 70 metres. Sample interval for sensors was 7-8 metres.

The survey was based out of Sudbury and completed on July 28, with complete coverage of the Platinex Shining Tree claim group. The calculated first vertical derivative of the magnetic field (Figure 11) and Anomaly Map from VLFEM survey (Figure 12) are provided from this work.

Z. Dvorak (2008) provided an interpretation of the Terraquest survey. The following are the highlights and a map (Figure 13) is included.

- A swarm of diabase dikes with azimuths of 160° to 170° is a prominent feature across the property with some 22 dikes evident forming narrow linear anomalies on Calculated Vertical Derivatives of Magnetic Field 'CVD'.
- A 110° fracture F1 associated with change in direction of diabase cuts across Cryderman Lake.
- Second fracture F2 is associated with well defined magnetic anomaly M1 associated with dunitic komatiite near south west corner of property. The area A is a reflection of high potassium and low magnetic values and has been mapped as a mafic to ultramafic intrusive. A hook like magnetic anomaly M2 and M3 is separated by 050° fracture F3 which trends from southwest of property to Michiwakenda Fault at Foisey Lake. Evidence for the fault is provided by both topography and magnetic data.
- Anomalies M6 and M7 are believed to reflect iron formations which are not magnetic everywhere. Area B has been mapped as Timiskaming Clastic metasediments which is close to the iron formation (M6).
- The CVD data near the Herrick deposit shows excellent correlation with bedrock features. The trachyte porphyry is defined by anomaly M8 and the veins are expressed as breaks in the magnetic patterns. Anomaly M4 which appears discontinuous from south of Hwy 560 appears to intersect the Herrick deposit at a low angle. Parallel anomaly M5 appears to lie just east of the known extent of the Herrick deposit.
- Anomaly M9 wraps an area of intermediate to mafic volcanics around the periphery of which several gold deposits including Ronda, Caswell and Bilmac are located. This feature may actually be terminated by the F3 fracture set which also appears to penetrate the Michiwakenda fault. Dvorak also pointed to the unique magnetic feature M2 centred in the NE corner of the McBride lease. It is now suggested that this may be related to the syenite intersected in drill hole MP-12-01 to the northeast.

- This magnetic anomaly coincides in part with VLF ORTHO anomaly V02 which appears to terminate at the Michiwakenda Lake fault. VLI is a linear VLF anomaly that appears to associate with an 'iron formation'.
- Dvorak draws attention to several of the VLF ORTHO anomalies that appear to represent bedrock features (Figures 2 and 13). He notes that the radiometric potassium anomalies appear to reflect positive areas but draws attention to an ORTHO anomaly coincident with a potassium anomaly on the west side of Caswell.

11.3 Ground Geophysical Surveys

Platinex has completed several focussed ground geophysical surveys on the property including walking magnetometer and pole dipole IP surveys. Numerous small blocks of EM and magnetic surveys have been done at various times and locations, by various interests, on the current Platinex Shining Tree claim block. No comprehensive ground geophysical surveys of the area have been documented. Compilation of the smaller surveys is ongoing, but will not provide complete coverage of the property.

JVX 2010a was commissioned to conduct an IP and magnetometer survey over the Herrick deposit covering 3.075km of IP resistivity and 6.818km of magnetometer surveys. The Herrick deposit was found to not be chargeable at surface but other parallel anomalies were found specifically to the east of the Herrick Central zone that warrant follow up work.

In 2011 JVX (2011a & b) conducted ground magnetometer surveys over Michiwakenda Lake and Perkins Lake covering 70 and 33.5 line km respectively. The line spacing was variable but approximately 12.5m and the readings were taken every second or at 2m intervals approximately coupled with GPS readings. The data enable accurate mapping of the diabase and trachyte and enable identification of structures.

11.4 Down Hole Geophysical Surveys

In 2010 Platinex commissioned a downhole IP survey by JVX (2010b). The survey was conducted from September 29 to October 3, 2010 and carried out on six holes HP09-33, HP10-43, HP10-44, HP10-45, HU89-6 and HU89-9. Results vary from hole to hole but one prominent anomaly is indicated in HP09-33 at the bottom of the hole and to the south. The zone is related to a chloritized zone in core which has not been assayed.

In 2011 DGI Geoscience was engaged to conduct downhole surveys of six holes on the Caswell prospect. A multi test process was used down the hole to provide structural orientation and also employing an IP probe and a natural gamma log probe to provide information on chargeability, resistivity and alteration (dominantly potassic).

12.0 GEOCHEMICAL INVESTIGATIONS

12.1 Regional Lake Sediment Survey

The Ontario Geological Survey and the Geological Survey of Canada (GSC) have performed a regional lake sediment and lake water survey in the Shining Tree area. Multi-element analyses of lake sediment and lake water samples from the Shining Tree area were released in 2001 report by Russell and Hamilton. The ICP-MS analytical technique was used for a wide variety of elements from both the sediments and lake water samples. The FA /ICP-MS technique was used for Au, Pd and Pt.

Two anomalous areas were defined in close proximity to, but not coincident with, the Shining Tree property. The "Area B" anomaly is comprised of 15 lakes from South Sandstrum Lake, about three kilometres east of the main block, to Jerry Lake in the northeast, about one kilometre south of the east extremity of the east block. Anomalous elements included the PGMs and some base metals. Most pertinent to this project is the indication that anomalous gold was present down-ice from the eastern portion of the Platinex Shining Tree property. Sites 277 and 382 returned concentrations of 5 to 7 ppb Au.

Another location, Site 279, adjacent to anomalous site 277 did not return anomalous gold in the OGS survey but had returned a value of 95 ppb Au in a GSC survey (Hornbrook and Friske, 1988). The presence of a gold occurrence on the western shore of the lake may, or may not negate the possibility of gold occurrences in the up-ice direction. It is also worth noting that there are no lakes between Jerry Lake and the West Montreal River, that is, in the down-ice direction from the south eastern part of the east portion of the Platinex property.

Johns (1999) showed the bedrock geology to be a northwest trending band of intermediate to felsic tuff breccia which is dissected by a gabbroic intrusion. A number of small Matachewan diabase intrusions have also been mapped. The geophysical data for the area shows a number of small, discontinuous magnetic highs which may represent additional intrusions without surface expression. Nearly all of the geochemical anomalies are within lakes entirely or partially underlain by the metavolcanic rocks. In addition, much of the area is covered with ice-contact glacial deposits. The materials which make up the ice-contact deposits are not necessarily sourced proximal to the area, and may also have some influence on some of the lake sediment results. The fact that the anomalies show some correlation to the underlying bedrock and are not scattered throughout the area is encouraging.

The second anomalous area, referred to as "Area 12: West Shining Tree Area", was located west of the Platinex property. Nine to seventeen lakes had elevated to anomalous Cu, Cd, Zn, +/- REE. One of the three most anomalous lakes (Site 1037) returned a sample analysis of 205.2 ppm Cu, 273 ppm Zn, 3.62 ppm Cd, 133 ppm Cr and highly anomalous REEs (Y and Be). The base metal anomalies indicate the associated volcanic stratigraphy could host base metal deposits and the OGS suggested the REE results may represent an area of hydrothermal alteration of the mafic to intermediate volcanic bedrock.

12.2 Lithochemical Studies

Petrographic work was done by the OGS, using samples collected by H.S. Oliver and G.W. Johns from 1997 to 1999, generally concentrated along the access provided by Highway 560. More than 380 samples were taken from the Keewatin assemblages, more than 70 from the Timiskaming volcanoclastics, and 50 from Timiskaming sediments. Trace element/immobile elements were along with medium and heavy rare earth element analysis, were used to classify and differentiate lithological units with an emphasis on differentiating mafic to ultramafic units.

12.3 On-Site Geochemical Surveys

12.3.1 Basal Till Sampling

In June of 2008, a reconnaissance till sampling program was initiated over the Platinex Shining Tree property. Subsequent till sampling has been carried out in 2010 and 2011. From 2008 through to June 2011 samples were taken from hand dug surface till pits. The sampling pattern took advantage of highway, logging road, trail and lake access and focused on the central, west and northern portions of the property. The property has moderate topographic relief and till is reasonably well exposed on hills and higher plateau areas with bedrock often within 1.5 metres of surface. Lower areas are covered with sandy outwash or organics greater than 1.5 metres thick and are not practically sampled with hand dug

Summary Report for Platinox Inc., Shining Tree, Ontario

By: James R. Trusler

pits. The later 2011 sampling was done using trails and a backhoe or excavator. There were a total of 307 samples taken but some of these were later rejected as silt, sand, gravel or organics.

The later 2011 sampling program was subject to a different process. First of all the samples were taken at approximately 50m intervals along trails or beside roads. Where possible a sample of bedrock was taken for identification and subsequent analysis. The samples were preanalysed and a smaller number were sent for processing by ODM.

The goal for each hand dug pit was to obtain 12-15 kg of screened sample, taken from beige, relatively non-oxidized till, close to or on the bedrock/till interface (basal till). Screening of large pebbles and cobbles larger than 6mm reduced shipping and handling weights and a rep sample of > 6mm pebbles for each sample was archived. The backhoe sampling achieved smaller average samples of 5 to 8 kg after screening.

Samples were shipped to Overburden Drilling Management (ODM) Limited for processing. All samples were processed for gold grain recovery and 12 samples were also processed for kimberlite indicator minerals. Gold grain recovery data is presented as the number of individual grains for each sample that fall into the shape categories of round, modified and pristine. Pristine grains have delicate features preserved, modified grains retain some form of the delicate features, and round grains exhibit no original features. These criteria enable an estimate of grain transport to be made in addition to defining gold dispersal trains based on numbers of grains.

A total of 446 till samples were processed by ODM. A total of 4,792 gold grains were recovered, of which 737 are modified and 595 are pristine. A total of 29 samples had greater than ten pristine plus modified gold grains.

Kimberlite indicator minerals other than chromites were present in six of the twelve samples processed.

Total counts were six chrome diopsides, one pyrope garnet and two kimberlitic olivines. High chromite counts occurred in most samples and were interpreted to be from ultramafic rocks south of Timmins (Averill, personal communications). No probe work on kimberlite indicator minerals was done.

Of the 446 till samples processed by ODM 209 samples contained 9 gold grains or more. These samples are clustered in two large areas (Figure 17) which define two of the more significant gold dispersion trains in the Abitibi gold region. Further 14 samples contain 50 or more gold grains including 3 that contain more than 100 grains.

The main gold dispersion train is elongated along a 060° trending axis which coincides with several noted geological and geophysical features. In addition some of the higher total gold grain counts and the higher pristine plus modified gold grain counts tend to cluster along the northern edge of the gold train and also about a circular feature coinciding with an East West trending XDS ORTHO anomaly west of Michiwakenda Fault and straddling Hwy 560. It is suggested from the drilling program that an alkalic stock is associated with much of this area. Also an ENE trending deformation zone identified by Johns (1996) crosses the Michiwakenda Lake Fault and coincides with the gold dispersion train. This deformation zone is believed to reflect the Larder Lake Break or a related splay (Figures 6, 17, 18).

The heavy mineral fractions from 199 of the till samples were subjected to whole rock analysis by ALS Chemex of Vancouver. Several parameters show anomalous results on the property. There is a correlation in general with the deformation zone and with NNW trending features such as the Michiwakenda Fault. Figures 19 to 22 display the variation in values for arsenic, sulphur, nickel and copper.

13.0 DRILLING

13.1 Historic Drilling

The most significant drilling completed to-date on the Shining Tree property was at the Herrick Deposit and to a lesser degree at the Churchill occurrence. The location of only one drill hole collar has been confirmed. Another eleven holes were completed by Unocal in 1989.

13.2 Platinex Drilling

From 2009 to 2011 Platinex completed 51 holes for 6181m on the Herrick deposit and strike extensions, 7 holes for 1070m on the Caswell occurrences in 2011. In 2012 Platinex drilled 5 holes for 870m to follow up coincident gold in till and IP anomalies in the Clark, Beilby and McBride areas. Due to budget limitations the logging and sampling of all five holes could not be completed and only selective work was done after a quick log of all five holes.

All 61 holes are shown on Figure 7 along with the surface geology as mapped by Unocal and later supplemented by Mehmet Spaho. Differences in interpretation still need to be reconciled, Cross Section 13+ 35N, is provided as Figure 14 without Platinex data added. The gold grades and intersection widths are provided on the section. Of particular note is the apparent continuity of the Central and East gold-bearing zones to depth. The drill hole plan Figure 7 and Longitudinal section Figure 15 are also shown.

14.0 DATA COMPILATION AND SYNTHESIS

14.1 Herrick Deposit

Complete paper records of Unocal drilling and channel sampling, including assay certificates, for the Herrick deposit were located and compiled into a digital database. Exploration records from work done by Herrick Gold Mines between 1918 and 1923, were compiled in 1935 by Grantland Gold Mines Ltd. and Erie Canadian Mines compiled a 300' level plan with sample locations. Collar location information was limited. Four drill holes were presented on one section with the distance of the collar denoted from either the shaft or vein. Northings were inferred for holes one and four from the presence of the shaft in the section in the first case, and a note on there being an orebody in hole four near the shaft. Hole 2 was located in the field by Unocal, and the location of hole 3 was inferred from a location on the Churchill and Connaught geology map (Carter 1980). Quartz-rich zones and geology are marked on the original Herrick Gold Mines sections which aid in correlating location, as well as confirming to some degree the location of the gold-bearing zones. The data for Herrick Gold Mines drill holes has not been placed in the database due to the lack of documentation; thus, these holes were not plotted on the sections or plan (Figures 7 and 14). However, approximated pierce points of the drill holes have been added to the longitudinal section of the Central Zone (Figure 15) so as to present the data on an approximate basis that adds some value to the overall interpretation. The drilling by Platinex was unable to confirm the reported tenor of the Herrick Gold Mine drill holes.

Channel sampling of the stripped portions of the Herrick vein structure have been done by several companies over the years, with the earliest documented in 1918 by Herrick Gold Mines. Results from this data are known only from a compilation done by mining engineer H.B. Hatch for Grantland Gold Limited in 1935. The Herrick channel sampling occurred in five areas identified as A, B, C, D and E along with calculated average values and widths. Lengths of channels average approximately 1.25m, with the majority between 1.0 and 1.5 metres. Several of these channels are evident in outcrop and are similar to channel samples pre-dating 1940 that were recently observed at the Caswell and Speed

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

power-stripped areas. Channels were hand moiled at a consistent depth and width (generally wider and shallower than saw channels) and give an impression that care was taken to representatively sample the structures but representing probably triple the volume of sample compared to saw channels.

Extensive sampling (201 channel samples, 35 composite grab/chips) was done by Unocal Canada Limited in 1989 after power stripping the area. This information is well documented in an exploration report by G. Robert Cluff, including assay certificates, and sampling maps. Locations and some of the grades and channel width information are provided on a representative cross section (Figure 14), and on the longitudinal section (Figure 15).

Additional sampling of the Herrick vein structure was completed by Fort Knox Resources as part of a more regional examination of the Shining Tree area. Forty-five channel samples were taken and documented by Doug Hunter in 1990. Sample locations and grade/width information are included on Figures 14 and 15.

Following is a summary of surface sampling work done to date on the Herrick vein structure:

- Three channel samples at the shaft area by Herrick Gold Mines reported gold values of 15.9 g/t over 1.25 metres, 24.9 g/t over 1.25 metres, and 16.9 g/t over 1.25 metres. Sampling by Unocal returned 8.8 g/t over 1.5 metres. Sampling by Fort Knox Gold Resources had yielded 13 g/t over 0.4 metres.
- North of the shaft, in what is now interpreted to be the Central Zone, Herrick Gold Mines reported 32 channel samples taken over a 50 metre strike length averaged 7.5 g/t over 1.25 metres, with the highest assay at 14.6 g/t and two channels returning Nil gold values. Unocal channel sampled 10 sections in this area, with best results of 14.7 g/t over 1.0 metres, 6.6 g/t over 1.0 metres, 9.8 g/t over 2.0 metres, and 7.9 g/t over 1.8 metres. Fort Knox cut three channels in this area with results of 3.31 g/t over 1.15 m, 9.49 g/t over 1.8m and 12.76 g/t over 1.75 m.
- Immediately south of the shaft the Central Zone continues. Herrick Gold Mines reported that 17 channel samples on this part of the Central Zone were taken over a 25 metre strike length and averaged 25.5 g/t over 1.15 metres. Thirteen samples returned assays over 15 g/t, with the highest assay 106.3 g/t and lowest assay 3.3 g/t. Channel sampling of this area by Unocal returned 56.5 g/t over 1.0m, 16.77 g/t over 0.5 metre, 4.8 g/t over 1.0 metre and 0.82 g/t over 1.0m. Visible gold was noted in outcrop. No channel sampling was done by Fort Knox in this area.
- South of the diabase and 100 metres south of the shaft, Herrick Gold Mines reported 28 channel samples taken over a 50 metre strike length of the vein, averaged 20 g/t over 1.3 metres. The highest gold assay was 200 g/t and the lowest assay was 4.0 g/t. Five samples assaying over 15 g/t. Unocal cut 12 sections of channel samples in this area, with the best gold assay results being 12.5 g/t over 1.5 metres, 3.9 g/t over 2.0 metres, and 5.7 g/t over 2.6 metres. Composite grab samples taken from the vein on the north part of this section returned gold values of 30 g/t, 20.41 g/t, 16.01 g/t, 2.54 g/t, 2.4 g/t, and 2.02 g/t. Fort Knox focused their sampling in this area with 7 channelled sections which returned 7.9 g/t over 0.5 metres, 5.5 g/t over 0.45 metres, 2.85 g/t over 1.65 metres, 4.0 g/t over 1.15 metres, 5.1 g/t over 1.25 metres, 5.0 g/t over 0.85 metres, and 5.56 g/t over 1.7 metres.
- Fifteen metres south of the current Unocal grid of 11+50N, Herrick Gold Mines reported 10 channel samples over a 15 metre strike length, averaging 7.2 g/t over 1.2 metres. The highest gold value was 11.3 g/t and the lowest gold value was 0.4 g/t. Unocal cut 5 channels in this area, with the best results being 4.9 g/t over 1.4 metres and 4.2 g/t over 1.0 metres. Four grab

samples from the area returned gold values of 25 g/t, 25.91 g/t, 0.75 g/t and 0.43 g/t. Fort Knox cut five channel sections in this area, which returned: 26.94 g/t over 0.88m, 9.02 g/t over 1.25m, 0.94 g/t over 1.0m, 4.90 g/t over 0.8 m, and 5.16 g/t over 1.3 m.

The only documented underground work was done by Herrick Gold Mines Limited between 1918 and 1923. Sampling of the top 25 metres of the shaft returned consistently high gold grades of 50 g/t across 1.5 metre widths (Figure 15). Eighteen samples were taken, with the highest gold value at 329 g/t and the lowest value at 6.4 g/t. Fourteen samples were greater than 15 g/t, eight samples greater than 30 g/t and three samples were greater than 100 g/t.

The only other documented underground assay results come from a level plan of the 300 foot level, redrawn by Erie Canadian Mines in 1935. The locations and grades have been indicated on Figure 15. The Central zone was mapped and sampled from a point approximately 6m south of the shaft, south approximately 60 metres to the contact with the diabase dyke. On the south side of the diabase dyke, a vein structure (possibly the West Zone based on drill section interpretation) is sampled for 30 metres, with the end of drift sample having a gold assay of 4.11 g/t over 0.33 metres. Samples were taken across either the back or face at approximately 1.5 metre intervals. Significant gaps in sampling occur where the structure was deemed to be too narrow or where no quartz veining was observed. Sample widths range from 0.2 to 1.7m across the shear/vein structure.

Four sections on the 300 foot level that have both continuous sampling and some of the better individual sample grades and widths are as follows:

- Central Zone: Centred on section 12+95, a 12.5 metre length of vein assayed 6.55 g/t over an average width of 0.78 metres; Centred on section 12+60, a 6.4 m length of vein assayed 9.62 g/t over an average width of 0.45 meters;
- West Zone: Centred on section 11+90, a 10.7 m length of vein assayed 9.87 g/t over an average width of 0.97 metres; Centred on section 11+75, a 17.3 m length of vein assayed 3.76 g/t over an average width of 3.76 metres.

Sampling of the drift was sporadic, resulting in sections of vein with good grade lacking continuous sampling. An example is a 3.4 metre long section of vein assaying 12.25 g/t over an average width of 0.7 metres. The sample plan denotes no sampling due to a lack of veining for a few metres both north and south of this section along the drift.

14.2 Caswell area

Previous work on the Caswell occurrences have been compiled and presented on Figure 8. A description of previous work was in section 8.2 herein.

14.3 Churchill area

The Unocal data from the 1989 channel sampling and drilling program have not been entered into the database. The bedrock was mapped by M. Spaho. The location and sampling results of underground development is unknown.

14.4 Speed Lake Area

No compilation has been done in this area. Some drill hole results just west of the property are available and should be added to a database along with all other information regarding the Gosselin

trend in Churchill and Asquith Townships. Recent till and airborne surveys completed by Platinex Inc. will add greatly to compilation work in this area.

15.0 SAMPLE COLLECTION, HANDLING AND PREPARATION

Exploration Work Pre-1989

No documents exist detailing the sampling methodology prior to 1989.

1989 Exploration Work done by Unocal (Churchill and Herrick Areas)

Channel samples and split diamond drill core were sent to Swastika Laboratories in Kirkland Lake, Ontario. Drill core was of NQ size and mechanically split, with one assay tag remaining in the core box. Core is stored at "Camp 560" located on the north side of highway 560, between the Bay Lumber Road and Houston Lake tourist camp.

Channel sampling was well done, and accurately mapped. Splitting/sampling of drill core appeared carefully and well done, with blocks and sample tags generally still intact. Wooden boxes are beginning to weather badly in a few cases, but are well labelled. It is perceived especially in view of drilling by Platinex that all of the core in the vicinity of the veins on Herrick should be systematically assayed.

1990 Exploration Work by Fort Knox (Herrick Area)

Channel samples were sent to Swastika Laboratories in Kirkland Lake, Ontario. Samples in the Herrick area were generally taken in areas of Unocal sampling, possibly as a direct check on previous results. Removal of sample material from channel cuts was acceptable.

2008 – 2011 Platinex Rock Samples

In 2008, Platinex took grab samples, chip samples and channel samples from various areas on the Shining Tree property as part of a due diligence program and a preliminary phase of exploration. Grab samples and chip samples were also taken of representative rock and historic muck piles by Platinex personnel.

Channels samples were taken from power-stripped areas, during and after the bedrock was mapped, using a gas powered masonry saw utilizing a water-cooled diamond impregnated blade. Sample lengths were recorded in assay books and samples were chipped from bedrock, placed in individual heavy duty plastic sample bags and labelled. Channel cutting and sampling work was performed by personnel from Katrine Exploration Services, Larder Lake, Ontario, under supervision from Platinex field geologists. Each sample received a pre-numbered assay tag from the assay book, and an aluminum tag labelled with the corresponding sample number was placed at the end of each channel cut.

Rock samples were gathered and transported to camp in plastic rice bags or plastic pails and subsequently sealed and delivered by Platinex personnel to the ALS Chemex prep lab in Sudbury.

2008 - 2011 Platinex Till Samples

(See section 12.3.1 Basal Till Sampling)

Till samples were sealed in plastic pails and shipped by Manitoulin Transport to Overburden Drilling Management in Nepean, Ontario.

The author is unaware of any drilling, sampling or recovery factor that could materially impact the accuracy and reliability of the results. To the extent known, the author is also unaware of any factors that may have resulted in sample biases.

16.0 SAMPLE ANALYSIS, QUALITY CONTROL, AND QUALITY ASSURANCE

Exploration Work Pre-1989

No documents exist detailing the analytical methodology, quality control or quality assurance protocols for work done on the property prior to 1989.

1989 Exploration Work done by Unocal (Churchill and Herrick Areas)

Gold analyses were done by Swastika Laboratories of Kirkland Lake, Ontario using 1 assay ton fire assays. Examination of the assay certificates indicates that check assays were done approximately every 10 samples and a second pulp with an additional check assay was done approximately every 25 samples.

Approximately 125 pulp and metallic assays were done on drill core samples from mineralized zones at the Herrick and Churchill to test for possible nugget effect (Cluff, 1990). The sample is crushed, pulverized and sieved to 100 mesh. Two analyses of the -100 mesh and one analysis of the +100 mesh are done by fire assay methods. The two -100 mesh values were averaged and combined with the +100 mesh value on a weight ratio basis to give a final calculated value.

Cluff estimated a 9% increase in gold values using pulp and metallic analysis, over the standard one assay ton analysis, but also noted from the results that nugget effect was not a problem (Cluff, 1989)(See following tables).

No blank, duplicate core, or assay standards are known to have been used. The location of pulp and reject material is unknown. The Report states that “this does not reduce the apparent validity of the results since there was excellent repeatability of check assays and the validation from samples collected by Bryant Groundwater Consulting was acceptable.”

1990 Exploration Work by Fort Knox (Herrick Area)

Fort Knox submitted 45 channel and grab samples to Swastika Laboratories of Kirkland Lake, Ontario, for analysis by fire assay. Eight samples had duplicate analyses. Assay results for the five samples with the highest concentrations of gold are provided in Table D4 of Appendix D. No other information is available.

2008 Platinex Rock Samples

Gold analyses of grab, chip and channel sampling done by Platinex Inc. in 2008 were done by ALS Chemex Laboratories in Vancouver, British Columbia. Initial crushing and pulverizing were done in an ALS Chemex prep lab in Sudbury, Ontario.

Sample preparation consisted of crushing of the entire sample to > 70% passing – 10 mesh (-2 mm). A 250 g riffle split was pulverized to >85% passing minus 75 um. One 50 gram aliquot of pulverized (pulp) material is fire assayed and finished with atomic absorption analysis to determine gold grade. Analyses greater than 10g/t would have a second fire assay performed with a gravimetric finish.

The screen fire assay method is used to detect the presence of coarse gold. Several samples from the Speed Lake, Caswell areas were submitted to ALS Chemex for this procedure. In this case 1000 grams of pulp material is sieved through a 100 micron dry screen to separate any coarse (+100 micron) material. Any +100 micron material remaining on the screen is dried, weighed and analysed in its entirety. The -75 micron fraction is homogenized and duplicate 50 gram sub-samples are analysed using fire assay. Both +75 and -75 micron fractions are weighted and reported together as a calculated total gold content.

Platinex Inc. relied on ALS Chemex internal standards and quality control procedures for the 2008 initial exploration work. Pulp and reject material for all samples sent to ALS Chemex have been shipped to Platinex and are currently stored in the company's Peterborough, Ontario core storage facility.

2008 - 2011 Platinex Till Samples

Overburden Drilling Management (ODM) of Nepean, Ontario processed the till samples taken from the Platinex property. A heavy mineral fraction was obtained from the 5-15 kg original sample using a shaking table and heavy liquids. Micro-panning was also employed on selected samples. Kimberlite indicator minerals gold grains and other heavy minerals were identified by ODM staff in size fractions between 0.25 and 2 mm under binocular microscopes. For gold grains, total counts, sizes, grain shape and calculated assay are tabulated for each sample. Geochemical analysis of the heavy mineral fractions have been completed for 199 samples out of 446 at ALS Chemex.

No duplicate samples have been submitted to ODM.

The sample preparation, security and analytical procedures employed by Platinex Inc. generally meet the industry best practices. The author is of the opinion that the work completed to date by Platinex is reliable.

17.0 DATA VALIDATION - 2008

17.1 Site Visit and Check Sampling

J.G. Bryant, visited the property with D. Jamieson on August 6-8, 2008, inspected three areas for evidence of prior work, independently collected and observed the collection of some outcrop chip samples from prior channel sample locations and outcrop and examined and sampled archived core from the Unocal Herrick drill program. Details of principal features, samples collected and analytical results are provided in the following sections according to showing or deposit name.

17.1.1 Herrick Deposit

- Most of the extensive cleared, stripped and washed areas of quartz, quartz-carbonate and quartz-carbonate-sulphide zones were well exposed.
- The shaft was observed to be partially blocked within several metres of surface.
- Abandoned equipment including a winch, boiler and stack were present in the overgrown brush southeast of the shaft.
- There was excellent correlation of the detailed, map-identified geology with observed features.
- Unocal channel sample locations and sample intervals were readily confirmed.
- Composite chip samples were collected from identified anomalous segments of Unocal channel sample. The samples were labelled and hand-delivered to the ALS Chemex lab in Sudbury for processing. Representative samples were analyzed by the ALS Chemex lab in Vancouver by the technique described in the previous section.
- Drill hole casings for collars of Unocal holes HU89-6, HU89-9, HU89-11, and HU89-12 were located in the field and tied in using hand held GPS units.

TABLE 2
VALIDATION SAMPLING RESULTS FOR HERRICK DEPOSIT – 2008

UNOCAL CHANNEL SAMPLE (1989) GOLD ASSAY RESULTS				BGC QUARTER SPLIT CORE (2008) GOLD ASSAY RESULTS			
Section	Sample No.	Interval (m)	Gold g/t	Sample No.	Length (m)	Gold g/t	
14+10N	7649	0.4	6.96/6.65	982	0.4	9.87	
13+72N	7606		7.2	983	0.4	7.58	
13+51N	6139	0.9	14.67/14.74	984	0.9	23.9	
12+90N	6127		16.77	See 12 +90 - 12 +93			
12+93N	6129		4.8	See 12 +90 - 12 +93			
12+90 - 12+93N	Composite chip sample from along quartz-Fe carbonate vein with sulphides.		None	Not sampled	986	3	15.4

HERRICK WASTE DUMP

Sample Description	Sample No.	Gold g/t
Grab Sample: Quartz and minor carbonate vein (suspected gold sheen on broken surface)	981	10.9
Composite of fine crushed and broken rock from waste pile	985	2.63

HERRICK WASTE DUMP

Sample Description	Sample No.	Gold g/t
Grab Sample: Quartz and minor carbonate vein (suspected gold sheen on broken surface)	981	10.9
Composite of fine crushed and broken rock from waste pile	985	2.63

Notes: BGC samples collected by G. Bryant, assays conducted by Swastika Laboratories; not an ISO accredited laboratory.

The author has not independently conducted validation tests to supplement data verification by Bryant and Jamieson. However, all of the subsequent work to that reviewed by Bryant and Jamieson has been supervised by independent qualified persons who have adopted appropriate sample handling and analysis procedures to ensure quality control at all times.

18.0 COMPETITOR ACTIVITY

Current and recent exploration and/or development activity in the Shining Tree area has been carried on by at least seven companies. The companies include Golden Harp Resources Inc. (Golden Harp), the Goldeye Exploration and Temex Resources Corp. (Goldeye/Temex) joint venture, URSA Major Minerals Incorporated, Creso Resources, Mineral Mountain Inc., Sarissa Resources (now Shining Tree Gold), Nirek Resources and RA Resources (now Golden Phoenix Resources).

Golden Harp has accumulated a large land holding east, north and northwest of the Platinex property. Their property, referred to as the Copper Hill Project, includes all but the southwest corner of Macmurchy Township and adjoining parts of Churchill, Natal, Knight and Tyrrell Townships. Although no gold deposits have been defined in exploration from 1990 to present, the technical report prepared by SRK Consulting in 2007 stated that the "project is of sufficient merit to recommend a two-stage exploration program that is designed to identify new targets while continuing exploration work on known gold occurrences" and that "the cost for the recommended work program are estimated at CDN \$1.8 million". Golden Harp optioned their property to Benton Resources in 2008 which the company vended the option after a limited drill program to Mineral Mountain Resources who have carried out extensive reverse circulation and diamond drilling programs since on the Cook deposit.

Goldeye/Temex have been active on their Juby North JV gold property located one township east of Macmurchy Township. A Phase 3 drilling program was recommended to start on the Juby Main Zone in late summer 2008. A June 24, 2008 press release indicated their recent program extended the gold-bearing zones and demonstrated the potential for not only significant strike lengths with the potential for additional gold discoveries. Long intervals of lower grade gold mineralization have been intersected and contain narrower intervals grading up to 5.43 g/t gold over 3.65 metres. Another intersection included 15.0 g/t gold over 2.15 metres. Temex in a recent press release revealed that the Juby deposit has been calculated to contain 1.84 million oz.

URSA has delimited a nickel-copper deposit in Fawcett Township, southeast of the Platinex property. The Indicated Resource was reported as 1.02 million tonnes grading 0.71% Ni, 0.36% Cu and there was an Inferred Resource of 1.49 million tonnes grading 0.67% Ni and 0.36% Cu at a cut-off value of 0.30% nickel equivalent. A preliminary technical and economic analysis and preliminary metallurgical test work has been completed. Environmental baseline studies were underway in 2007. The project was evaluated for a 1,000 tonne/day open pit operation with haulage and processing at URSA's proposed Shakespear Mill in the Sudbury area.

Creso Resources Inc. Has assembled a large land package surrounding the Platinex Shining Tree property and including the Minto and Tyrinite deposits. Several drill holes have been completed on both deposits including one spectacular result drilling down the Minto breccias pipe which returned an intersection of 82.5m of 13.3 g/t Au (MC-09-01).

Nirek Resources Inc. has an 18 claim property that coincides with the Gosselin quartz veins in Asquith and Churchill Townships immediately west of the Platinex Shining Tree property. The Gosselin vein has a 2.0 kilometre strike length and trends north 15 degrees west. The association is gold-bearing quartz veins in altered pillow lava, rusty weathering Fe-Mg-calcium carbonate, and cross-cutting felsite, or rhyolite and granitic porphyry. Some evaluation of green carbonate zones with lower grade Au concentrations has also occurred. It was also noted that a competitor had performed extensive washing, trenching and sampling on a long quartz and quartz-carbonate vein between the Speed Lake showing and Highway 560.

RA Resources (now Golden Phoenix Resources) has performed work on claims adjacent to the Speed Lake showing in the southwest part of the main claim block.

19.0 MINERAL PROCESSING AND METALLURGICAL TESTING

The Shining Tree project is at an early exploration stage and there is no need for mineral processing or metallurgical testing.

20.0 MINERAL RESOURCE AND MINERAL RESERVE ESTIMATES

Evaluation of the Herrick deposit is at an early stage with the result that there is insufficient information at this time to declare the Herrick deposit as a mineral resource. However, the author has provided calculations based on available data which is either qualified and/or has been obtained by independent qualified persons. Efforts will be made in the subsequent sampling programs to expand the grade and continuity of the gold-bearing zones as originally investigated. Positive results would result in initiation of a mineral resource estimate.

The Company SEDAR-filed its NI 43-101 technical report “the Report” dated October 2, 2008, by J.G. Bryant and D. Jamieson which examines the Herrick gold deposit on Shining Tree property and seven other known gold prospects, some of which have been explored underground. The Report qualifies the sampling and drilling work by Unocal (1989) and Fort Knox (1990) compliant to NI 43-101 standards. This report draws on the previous Report and provides an update based on results of subsequent exploration.

The Report states that, “Judging from evidence of previous results on the Herrick deposit there is a potential to outline a small commercial gold deposit.”

An internal valuation of the Herrick deposit completed in May, 2012 by the author estimated potential gold content ranging from 111,000 oz to 172,000 oz Au within the defined vein structures to a depth of 300m. The estimate is based on a minimum of 1,628,000 tonnes grading 2.11g/tonne to a maximum of 2,554,000 tonnes grading 2.10g/tonne Au. The estimate is derived from 3 narrow sub-parallel quartz vein breccias and an enclosing mineralized shear zone within a 370m strike length, to a 300m depth and within a 80m wide block. A cut off grade of 0.5g/tonne was used in the calculations.

The Central zone of the Herrick deposit outcrops for most of its length and is a continuously readily definable body. The valuation also identified thicker gold zones between intersecting or bifurcating veins in some holes (e.g. hole HP10-44 to the southwest) and in outcrops which are not included in the calculations. These structures plunge near vertically providing a prospective target for deeper exploration. There is an undefined potential for a southwesterly plunging thickened enriched section that could contain from a depth of 300m to 1500m some 14.6 million tonnes at a grade ranging from 2.0 to 5.0 g/tonne Au for a gold content of 1.0 to 2.4 million ounces. The model is based on the mentioned structure, comparison to the West Timmins deposit and statistical grade estimates by the author. The potential quantity and grade is conceptual in nature. There has been insufficient exploration to define a mineral resource and it is uncertain if further exploration will result in the delineation of a mineral resource. The internal valuation was conducted by the author, who is a qualified person but an insider of the issuer and therefore according to the ethical standards set forth in the National Instrument 43-101 no qualification of resources is permitted. More sampling of existing core and stripping of identified thicker mineralized zones where these project to surface, is warranted and could increase the identified potential within the exploration envelope and lead to a resource estimation by an independent, qualified person .

21.0 DISCUSSION

21.1 Historic Context

The Shining Tree gold camp underwent an initial surge of development from 1912 to 1925, and then again from 1933 to 1939. Since that time there has not been extensive exploration work done in the area. Unocal Canada Ltd. (Herrick) and Strike Minerals (Ronda) drill programs were the largest programs known by Bryant and Jamieson (2008) to be undertaken in recent times, and were each limited to a few months or less of diamond drilling.

The recent drilling campaigns by Platinex on the Herrick deposit and the property wide till studies are the most comprehensive exploration programs to date.

It is important to understand the history of a mineral-rich area, and attempt to understand the reasons for past development failures. Exploration reports written about the Shining Tree area often refer to the early days of the camp as having been less than orderly. For example, exploration campaigns consisted of “mini-rushes based on findings of spectacular gold occurrences...ending in investor losses” (Tindale 1989). H.C. Laird stated in his examination of the area in 1934” its career (Shining Tree gold camp) has been a hectic one from the point of view of mining development. In the early days of the camp the discovery of much spectacular high-grade gold ore, particularly in the Wasapika section, led to a period of extravagant financing and fanciful development. The inevitable result of this over-expansion was chaos, litigation, and a general loss of public confidence, a blow from which this camp has not yet fully recovered. At present attempts are being made to correct this condition by what is believed to be a sincere mining effort on the part of certain interests, particularly those holding properties in the Wasapika section).

Although the prospectors did a good job of finding surface showings, some or most of the subsequent exploration work may not have been systematic or of high quality. Interest in the area resumed with higher gold prices in the 1930's, however labour shortages due to the war caused the mining projects in the area to close.

The opening of claims for staking due to lapsing leases and patents in the last 20 years, as well as the general lack of activity in the area, provided an opportunity for Robert Macgregor to assemble a large land position that includes many of the key historic gold prospects in the Shining Tree gold camp. The property is large enough to enable more sophisticated regional exploration techniques such airborne geophysics, and till geochemistry/gold grain analysis to be employed. As a result an improved understanding of the geology and the geological control for gold mineralization could lead to lateral and depth extensions to existing deposits and occurrences and to the generation of new gold targets.

21.2 Spatial Context

Some of the key gold-bearing veins on the Platinex Inc. Shining Tree property cluster near the Churchill and Macmurchy township boundary north and south of highway 560. The Caswell vein system is part of a larger trend that includes four non-Platinex properties to the southeast that have undergone limited historic underground development, namely the Bilmac, Atlas, Bennett and Kingston properties. In addition, the Herrick vein system aligns along this same northwest trend, despite being separated by the Michiwakenda Lake Fault, which displaces stratigraphy for several kilometres along strike and an unknown vertical distance. Just to the west of the Caswell trend, the Ribble vein and Foisey vein on the Ronda property are north-south veins sub-parallel to the Michiwakenda Lake Fault and proximal to it (within 100 metres in the case of the Foisey vein). Similarly, the north-south Herrick vein structure, one kilometre to the north, shows the same relationship to the Michiwackenda Lake Fault, albeit on the opposite side of the fault.

21.3 Caswell area

The area has been subjected to a large number of relatively short exploration programs (except for the main underground development by Canadian Champion Mines) by a large number of companies, and compilation/interpretation suffers from undocumented or misplaced data. The successes of some early drill programs do not appear to have been duplicated by subsequent programs, although that may be due to the limited available documentation. Nevertheless, high grade gold values have been repeated by several workers in a few locations.

Review of compilation maps and the mapping and sampling of the area power-stripped by Platinex in 2008 confirm that the Caswell area has undergone complex strain and a prolific development of quartz veins hosted by narrow shear zones. Gold grades between 4 and 12 g/t were obtained across sub-metre intervals in channel samples across the No. 1 and related shear/vein structures exposed by the power-stripping. Most channel samples returned anomalous gold values.

Although mapping and sampling of the Caswell vein system on surface and underground has taken place in the past, there is no reliable documentation of these results.

Despite the negative aspects of Caswell data e.g. the historic very high grade mineralization which does not appear to be laterally continuous or in some cases repeatable, the Platinex work did demonstrate some positive characteristics and potential for the Caswell mineralization.

There is a proliferation of shear hosted ENE veining which intersect the historic apparently formational Evelyn and Saville veins. The ENE veins demonstrate a correlation of silver – tellurium and low grade gold mineralization throughout. Cutting (2012 personal communication) noted that the strong quartz flooding alteration in which he would expect to see potentially commercial gold grades in the Platinex stripped areas was limited to strike lengths of a few metres and widths of no more than three metres.

If this can be put in the context of the containing shear structure and/or it is related to an alkali-syenite system such as is postulated for the Minto pipe then an exploration approach can be established with a potential for success based in structurally controlled vertically oriented mineralization. There is a need for careful mapping of the shear hosted veins.

21.4 Herrick Deposit

The Herrick vein received relatively systematic early exploration from the time of discovery in 1918 until 1923 when the underground development program ceased. The project was dormant until 1989, except for undocumented further surface sampling in 1940. It should be noted that much of the current documentation of the early underground development work is derived from reports and examinations of the property data during the 1930's. The 1989 Unocal exploration program, the Herrick and Churchill areas, was a well documented and systematic exploration program that provided an excellent framework to compile early exploration results and develop future exploration programs.

The Herrick vein structure appears to be an extensive gold-mineralized system. Although the results of Unocal's work did not meet their threshold for continued expenditures, further work is needed to determine the extent of the local higher grade gold mineralization indicated, as well as the extent of wider zones of lower grade gold mineralization intersected by Unocal. The author believes that the Herrick gold deposit is a high priority exploration target that may be up-graded to a gold resource with additional stripping, channel sampling and additional sampling of available core.

Work by Platinex on the Herrick deposit has demonstrated a potential for a thick possibly good grade deposit plunging to the southwest at depth which could contain up to 2.4 million ounces. More modest mineralization is evident on surface, but the deposit might be larger than perceived due to ballooning

mineralization between vein splays. This needs to be tested by more sampling of existing core and stripping on surface.

There is also evidence that another east zone occurs within hole HP09-33 which was not assayed towards the end of the hole. There is ample potential for such zones between the Herrick Central zone and the Michiwakenda Lake Fault.

21.5 Churchill area

In the Churchill area, gold mineralization outlined by Unocal's 1989 exploration programs did not confirm earlier (1930's) work by the Churchill Mining and Milling Company. There is little documentation of the underground workings or assay results from the Churchill Mining and Milling Company's work. Some of the only information comes from visits by Ontario Department of Mines geologist H.C. Laird in 1934 and 1935. Laird had noted visible gold only in vein No. 1 and the west end of vein No. 3, and reported that company channel sampling at the time averaged 27.4 g/t over 1.25 metres on the No. 3 vein and 29.4 g/t over 1.25 metres on the No. 1 vein.

A partial explanation for the lack of confirmation of this early work by Unocal's exploration program lies in the fact that Unocal surface work did not examine the areas noted by Laird that is the No. 1 vein or the portion of the No. 3 vein west of the shaft. The four diamond drill holes that Unocal drilled in 1989 did not test beneath the No. 3 vein directly west of the shaft. Two holes were designed to test directly beneath No. 1 vein, however, one hole, CU-89-16, had to be abandoned after encountering underground workings. Hole CU-89-17 encountered altered trachytic porphyry with anomalous gold values that could correlate with the No. 1 vein. Hole CU-89-19 tested for the strike extension of the No. 1 vein beneath a large swamp, 100 metres to the east of the showings and also intersected the altered trachytic porphyry with anomalous gold values. The trachytic rocks encountered on the Churchill property are noted to be identical to those observed two kilometres to the north on the Herrick property.

An important indicator of base metal potential in the Churchill area is the presence of massive pyrite mineralization within cherty exhalite rocks. These rocks are along strike from regionally mapped sulphide iron formation, but magnetic surveys by Unocal show low magnetic response along this trend, indicating the absence of magnetite or pyrrhotite. The presence of sphalerite with the strong pyrite mineralization is suspected but not confirmed by Cluff, and no base metal analyses are provided.

The Churchill area requires additional examination due to its proximity to the Herrick gold deposit (see Figure 6) and to the lack of systematic exploration of the No. 1 and No. 3 veins.

21.6 Ribble vein (Ronda)

The depth extension of the northern portion of the Ribble vein may trend onto the Platinex Shining Tree claims at approximately 300 meters vertical depth. Testing of the vein below the depth of past production was done by Strike Minerals in 1996 and intersected sporadic high-grade gold values. This same mineralized level is projected to strike northward onto Platinex claims. The closest holes from the Strike Minerals 1996 drilling program are located about 60 metres south of the Platinex property. The Ribble vein depth extension is a high priority target for evaluation by deep drilling (400 to 500 metre long drill holes) on the Platinex ground.

22.0 CONCLUSIONS

The presence of three past-producing gold deposits and numerous gold prospects with underground workings in the Shining Tree gold camp underlines the significant potential for developing commercial gold production in this area.

In the context of regional geology and economic potential the Shining Tree property is part of the Abitibi gold belt and appears to be on a major gold associated feature the Cadillac Larder Lake Break. It is midway between major gold deposits (Cote Lake – IAMGOLD and Young Davidson – AURICO) associated with felsic intrusives on the same break. It is also on the same north-south structure as the West Timmins Mine of Lake Shore Gold. As with these three deposits the Shining Tree property has now demonstrated an association with an alkali (syenite) intrusive body. The intersections of major north-south faults (e.g. the Montreal River Fault) with gold associated features such as the Cadillac Larder Lake Break is a known harbinger of major gold and VMS deposits.

The Herrick deposit has a demonstrated potential to contain a significant gold resource. A program to strip and sample the veins at surface and thoroughly sample existing core is recommended.

Till studies on the Shining Tree property have demonstrated the existence of a regionally significant gold dispersion train which is believed sourced by several deposits possibly more significant than any known gold deposit on the property. Comparison of gold in till in Shining Tree to Timmins or Matachewan shows that Shining Tree has a high potential to host multiple commercial gold deposits (Figure 16 and 17).

Drilling of IP anomalies associated with a few of the high gold in till targets in 2012 intersected strong mineralization in four holes and an altered syenite in the fifth hole containing anomalous gold. Due to budget limitations most of the core could not be logged, sampled and assayed.

Much more of the property remains to be tested.

A two-phase exploration program is proposed to step up the evaluation of existing and anticipated gold-bearing targets during the balance of 2012 and in 2013. The proposed work program should include more core sampling, stripping and channel sampling, then deeper diamond drilling of the gold zones at the Herrick deposit and testing for the depth continuity of the Ronda gold-bearing zone beneath the Platinex property. Additional detailed surface prospecting, surface stripping and channel sampling, line cutting, pole-dipole IP, soil gas surveys and diamond drilling are also recommended. The proposed budget for Phase I is \$1,554,500, and the budget for Phase II is \$3,272,550 and the proposed work program for phase I is tabulated (Tables 4, 5 and 6).

Proposed drilling in Phase 1 totals 5,000 m and in Phase II 15,000m.

On the Platinex Shining Tree property, the Herrick deposit represents a high priority exploration target that could be upgraded to a gold resource with additional work.

High gold grain counts in till sampling indicates potential for new gold targets. While historic work indicates potential to discover deeper gold mineralization at the Ribble, Churchill and Caswell areas.

23.0 RECOMMENDATIONS

It is recommended that a phase I program totalling \$1,570,082 be carried out (Tables 4 and 6). Initial work is to involve completions of logging and sampling of the 2012 drill core then stripping and sampling selected areas over the Herrick deposit and much more sampling of the Herrick core.

Several areas where IP anomalies were found in combination with gold in till high counts may be stripped and sampled for initial evaluation. This is recommended.

Line cutting of 12 km is recommended to fill in the lines cut and surveyed in early 2012. IP, soil gas hydrocarbon sampling and magnetometer surveys are recommended with prospecting and geological mapping.

Summary Report for Platinox Inc., Shining Tree, Ontario

By: James R. Trusler

More follow up stripping and channel sampling and drilling is also recommended with an emphasis on using a portion of the drilling to follow up on Herrick exploration with one hole testing further east than hole HP09-33 and two holes testing Herrick beneath previous drilling.

**TABLE 4
PROPOSED PHASE I BUDGET
PLATINEX INC. SHINING TREE PROJECT, ONTARIO**

Phase & Tasks	Quantity	Units	Unit Cost	Proposed Budget
PHASE I				
-Qualifying report	1	1	\$ 25,000	\$ 25,000
-Project management & supervision	20	person days	\$ 800	\$ 16,000
-Mapping	15	days	\$ 750	\$ 11,250
Prospecting, core logging & splitting:				
-Project geologist	90	days	\$ 600	\$ 54,000
-Assistant	100	days	\$ 200	\$ 20,000
-Overburden trenching	150	hours	\$ 150	\$ 22,500
-Overburden stripping, washing, channel cutting and sampling	200	hours	\$ 150	\$ 30,000
-SGH	300	samples	\$ 50	\$ 15,000
-Line cutting	12	line km	\$ 750	\$ 9,000
-Ground geophysical surveys				
-Magnetometer	21	line km	\$ 200	\$ 4,200
-IP	12	line km	\$ 2,500	\$ 30,000
-Down Hole IP	10	holes	\$ 1,500	\$ 15,000
-Sample analysis				
-till samples (including collection costs)	100	samples	\$ 100	\$ 10,000
-rock and core	6,000	samples	\$ 27	\$ 162,000
-Drilling - core holes (20-25 holes)	5,000	metres	\$ 101	\$ 505,000
-Travel	10,000	km	\$ 0.55	\$ 5,500
-Equipment rental and field supplies			\$ -	\$ 25,500
-Meals & accommodations	150	person days	\$ 170	\$ 25,500
-Data compilation & drafting	32	days	\$ 200	\$ 6,400
-Reports	25	days	\$ 600	\$ 15,000
-Contingency	15%			\$ 151,028
Subtotal				\$1,157,878
Administration (20%)				\$ 231,576
HST (13%)				\$ 180,629
TOTAL				\$1,570,082

**TABLE 5
PROPOSED PHASE II BUDGET
PLATINEX INC. SHINING TREE PROJECT, ONTARIO**

Phase & Tasks	Quantity	Units	Unit Cost	Proposed Budget
PHASE II				
Advance Royalty payment April 2013			\$ 10,000	\$ 10,000
-Project management & supervision	30	person days	\$ 800	\$ 24,000
-Project plans, map, prospect, till sample, supervise drillers, log & split core	0	0	\$ -	\$ -
-Project geologist	160	days	\$ 600	\$ 96,000
-Assistant	160	days	\$ 200	\$ 32,000
-Overburden trenching	40	hours	\$ 150	\$ 6,000
-Overburden stripping, washing, channel cutting and sampling	70	hours	\$ 150	\$ 10,500
-Airborne geophysical survey				
-Consultant geophysicist				
-Line cutting	4	line km	\$ 750	\$ 3,000
-Ground geophysical surveys				
-Magnetometer	4	line km	\$ 200	\$ 800
-IP	4	line km	\$ 3,000	\$ 12,000
-Down Hole IP (10 holes)				\$ 20,000
-Sample analysis				\$ -
-till samples (including collection costs)	100	samples	\$ 100	\$ 10,000
-rock and core	6,000	samples	\$ 27	\$ 162,000
-Drilling - core (50 holes)	15,000	metres	\$ 101	\$1,515,000
- 10 holes	300	metres	\$ 300	\$ 90,000
-Travel	20,000	km	\$ 0.55	\$ 11,000
-Equipment rental and field supplies, including core racks	0	0	\$ -	\$ 35,000
-Meals & accommodations	180	days	\$ 170	\$ 30,600
-Data compilation & drafting	40	days	\$ 200	\$ 8,000
-Reports	40	days	\$ 600	\$ 24,000
-Contingency	15%			\$ 313,485
Subtotal				\$2,413,385
Administration (20%)				\$ 482,677
HST (13%)				\$ 376,488
TOTAL				\$3,272,550

TABLE 6: PROPOSED WORK PROGRAM PHASE I

<p>Complete logging and sampling 870m core:</p> <ul style="list-style-type: none"> -400 samples for gold -10 days geologist plus assistant
<p>Build core racks on lease near building for 15,000m core:</p> <ul style="list-style-type: none"> -set up house for viewing /storing core -consolidate core to that location -8 man days for 2 assistants
<p>Review core logs and core on Herrick:</p> <ul style="list-style-type: none"> -1500 core samples - 30 man days sampling - geologist days setting out samples and reviewing
<p>Line cutting:</p> <ul style="list-style-type: none"> - 10km fill in lines Beilby, Ribble Lake to Cryderman Lake and Clarke showing -2km line cutting on Herrick west to high till reading -Fix line numbering on winter grid 3 man days
<p>IP 12km; 2km on Beilby 2km on Herrick</p> <ul style="list-style-type: none"> -magnetometer 21km continuous read out
<p>Till sampling to follow up high till value locations:</p> <ul style="list-style-type: none"> -hand dug pits plus backhoe in some locations
<p>SGH sampling up ice from high gold in till and as check over IP anomalies plus over Herrick and to east:</p> <ul style="list-style-type: none"> -100m line spacing and 50m sample spacing with later fill in to 25m for 220 samples. -on Herrick and to east for 400m in 4 lines samples at 25m spacing for 80 samples -8 man days sample collection
<p>Geological mapping:</p> <p>of all lines, along highway and certain shorelines plus mapping of Herrick and other showings once stripped:</p> <ul style="list-style-type: none"> -15 days geologist and assistant -objectives to define evidence of major east west zone of high strain, provide lithological guidelines and map critical perceived ore controlling features including the alkalic (syenite) pluton
<p>Stripping, channel sampling:</p> <ul style="list-style-type: none"> -Herrick - expose splays and attempt to expose surface trace of West Zone -50 hours stripping, washing, channel sampling -50 hours overburden trenching
<p>Follow up IP, SGH and till results:</p> <ul style="list-style-type: none"> -150 hours stripping, trenching plus 100 hours overburden trenching -allow for 500 rock samples total including channels
<p>Reconcile geological interpretation with production of new plans and sections for Herrick:</p> <ul style="list-style-type: none"> -10 man days -2 geologist days
<p>Drilling:</p> <ul style="list-style-type: none"> - two deeper holes on Herrick, one or two holes to test down hole IP anomaly and any SGH anomaly. 1200m; 600 samples -one hole to test down dip of Ribble vein on Platinex property 400m, 200 samples - 20 holes plus or minus for 3400m to follow up anomalous indications of SGH, IP, stripping and trenching, 1800 samples
<p>Down Hole IP:</p> <ul style="list-style-type: none"> -10 holes tested depending on results of drilling; different IP systems are possible
<p>Qualifying Report:</p> <ul style="list-style-type: none"> -by independent geologist to include resource valuation on Herrick

24.0 REFERENCES

Ayer, J.A. 2000.

The Abitibi greenstone belt: a program overview; in Summary of Field Work and Other Activities 2000, Ontario Geological Survey, Open File Report 6032 p 6-1 to 6-14.

Ayer, J.A. 1999.

Geological compilation of the Abitibi greenstone belt in Ontario; toward a revised stratigraphy based on compilation and new geochronology results; Summary of Field Work and Other Activities, Ontario Geological Survey, Miscellaneous Paper 169, p.14-24.

Ayer, J.A., Berger, B. Johns, G.W., Trowell, N. Born, P. and Mueller, W.U. 1999.

Late Archean rock types and controls on gold mineralization in the southern Abitibi greenstone belt of Ontario; GAC.MAC Joint Annual Meeting, Sudbury 1999, Field Trip B3 Guidebook, 73p.

Ayres, L.D., Thurston P.C., Card, K.D. and Weber W. (eds.). 1985.

Evolution of Archean supracrustal sequences. Geological Association of Canada, Special Paper 28, p.343-375.

Carter, M.W. 1977.

Geology of Fawcett and Leonard Townships, districts of Sudbury and Timiskaming; Ontario Division of Mines, GR146, 50p.

Carter, M.W. 1980.

Geology of Connaught and Churchill Townships, District of Sudbury; Ontario Geological Survey Report 190, 81 p.

Carter, M.W. 1987.

Geology of the Shining Tree Area, districts of Sudbury and Timiskaming; Ontario Geological Survey, Report 240, 48p.

Casaceli, R., 2011

Personal Communication - discussion respecting gold-alkaline mineral systems.

Clark, R. 1989.

1989 Assessment Report Geological and Geophysical Surveys Churchill Project, Churchill Township, Ontario, NTS41P11. Assessment Report 41PSE0039.

Cluff, G.R. 1990.

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

Exploration Activities – 1989 Herrick and Churchill Properties, Churchill Township, Ontario. Unocal Canada Limited.

Barries, Charles, 2008.

Operations Report for Platinex Inc.; High Resolution TM-sensor Magnetic, KDs VLF-EM & Radiometric Airborne Survey; Shining Tree Project, Sudbury, Ontario; Terraquest Ltd. Report B-281.

Dube, B. and Gosselin, P. 2005.

Greenstone-Hosted Quartz-Carbonate Vein Deposits (Orogenic, Mesothermal, Lode Gold, Shear-Zone-Related Quartz-Carbonate or Gold-Only Deposits) in Geological Survey of Canada Open File Report

Dvorak, Zbynek, 2008.

Report on Preliminary of Airborne Geophysical Survey in the Shining Tree Area – An Interpretation of fixed wing airborne survey conducted by Terraquest Ltd for Platinex Inc. June 24-27, 2008.

Edgar, B. 2001.

Caswell Property Compilation/ Highlights and Property visit. Practical Exploration and Development Corporation Assessment Report.

Finley, F.L. 1926.

Wasapika Section, West Shining Tree Gold Area, District of Sudbury; Ontario Dept. Mines, Vol.35, pt.6, p.83-96 (published 1927).

Franko, S 2011.

Prospecting Report on active claim 4252127 Macmurchy Township (G-0988) Larder Lake Division 80 Ontario, Canada

Gordon, J.B. et al. 1979.

Gold Deposits of Ontario, Part 2: Part of District of Cochrane, Districts of Muskoka, Nipissing, Parry Sound, Timiskaming and Counties of Southern Ontario; Ontario Geological Survey, Mineral Deposits Circular 18.

Hopkins P.E. 1920.

West Shining Tree Gold Area; Ontario Dept. Mines, Vol.29, pt.3, p.28-52. Accompanied by Map No.29a, scale 1 inch to 1/4 mile.

Jamieson, D. 2009.

Diamond Drilling Assessment Report on the Herrick Gold Deposit, Churchill Township, Larder Lake Mining Division, Ontario, Canada.

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

Jamieson, D. 2010.

Till Sampling on the Shining Tree Project for Platinex Inc. Churchill, Asquith and Macmurchy Townships, Ontario, Canada.

Jamieson, D. 2011.

Diamond Drilling Assessment Report on the Herrick Gold Deposit, Churchill Township, Lader Lake Mining Division, Ontario, Canada

Johns, G.W. 1996.

Reappraisal of the Geology of the Shining Tree Area, districts of Sudbury and Timiskaming; Summary of Field Work and Other Activities 1997, Ontario Geological Survey, Miscellaneous Paper 166, p.13-15.

Johns, G.W. 1997.

Reappraisal of the geology of the Shining Tree area, districts of Sudbury and Timiskaming. Summary of Field Work and Other Activities 1997, Ontario Geological Survey, Miscellaneous Paper 168,p.26-29.

Johns, G.W and Amelin, Y. 1999.

Reappraisal of the geology of the Shining Tree area (East Part), districts of Sudbury and Timiskaming; Summary of Field Work and Other Activities 1999, Ontario Geological Survey, Miscellaneous Paper 169, p.43-50.

Johns, G.W. 1999a.

Reappraisal of the geology of the Shining Tree area (West Part), District of Sudbury; Summary of Field Work and Other Activities 1999, Ontario Geological Survey, Open File Report 6000, p.6.1-6.7.

Johns, G.W. 1999b.

Precambrian geology, Shining Tree area (east half); Ontario Geological Survey, Preliminary Map P.3389.

Johns, G.W. 2000.

Precambrian geology, Shining Tree area (west half); Ontario Geological Survey, Preliminary Map P.3420.

JVX 2010, Report of Borehole Induced Polarized Surveys Herrick Grid, Shining Tree project, Churchill Township, Northern, Ontario for Platinex Inc.; Ref No. 10-73 JVX Geophysical Surveys and Consulting.

JVX and Jamieson, D. 2010.

Report on Line Cutting, IP/Resistivity and Magnetic Surveys, Herrick Grid, Shining Tree Project, Ontario, Platinex Inc. Ref. 10-27.

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

JVX 2010a, Report on Line Cutting, IP / resistivity and Magnetic Surveys. Herrick Grid, Shining Tree Project, Ontario, Platinex Inc. Ref. 10-27.

JVX 2010b, Report on Borehole Induced Polarization Surveys, Herrick Grid, Shining Tree Project, Churchill Township, Northern Ontario, Ref 10-73.

JVX 2011a, Report on Magnetometer Surveys, Herrick and Caswell Grids, Shining Tree Project, Churchill Township, Northern Ontario, Platinex Inc. Ref 11-13

JVX 2011b, Report on Detailed Ground Magnetic Surveys, Perkins Lake Area, Shining Tree Project, Churchill Township, Northeastern Ontario, Platinex Inc. and Creso Exploration Inc. Ref 11-22.

Laird, H.C. 1934.

Geology of the Makwa-Churchill Area; Ontario Dept. Mines, Vol. 43, pt. 3, p.37-80 (published 1935). Accompanied by Map No.43c, scale 1 inch to 1 mile.

Laird, H.C. 1935.

Recent Developments in the Swayze and West Shining Tree Areas; Ontario Dept. Mines, Vol.44, pt.7, p.38-47 (published 1936).

MacGregor, R. 2008.

Report on Trench Geology and Channel Sampling Churchill and Macmurchy Townships, Ontario

McCannell, J.D. 1975.

Tri-Bridge Consolidated Gold Mines Limited, Asquith and Churchill Townships, Shining Tree Area, Ontario. Report on Geological Mapping and Diamond Drilling; Assessment file 41P11SW0025.

Newbury, M.2011

Personal communications, Report for New Bedford Explorations by Thomas Skimming & Associates 1975.

Oliver, H.S., Hughes, D.J., Hall, R.P. and Johns, G.W. 1999a.

Preliminary Geochemistry of Metavolcanic rocks of the Shining Tree Area; Abitibi Sub-province, Ontario; Summary of Field Work and Other Activities 1998, Ontario Geological Survey, Miscellaneous Paper 169, p.51-58.

Oliver, H.S., Hughes, D.J., Hall, R.P. and Johns, G.W. 1999b.

The Mafic And Ultramafic Volcanics Of The Shining Tree Greenstone Belt, Northeastern Ontario, Canada; Summary of Field Work and Other Activities 1999, Ontario Geological Survey, Open File Report 6000, p.11.1-11.12.

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

Oliver, H.S., Johns, G.W., Thurston, P.C., Hughes, D.J. and Hall R.P. 1998.

Ontario Geological Survey, Preliminary geochemistry of the Shining Tree area; Abitibi Sub-province, Ontario; in Summary of Field Work and Other Activities 1998, Miscellaneous Paper 169, p.51-58.

Russell, D.F and Hamilton, S.M. 2001.

Ontario Geological Survey, Lake Sediment and Water Data for the Shining Tree Area, Miscellaneous Release – Data 81.

Sinclair, D.G., Reeley, B.C., Cooper, D.F., Weir, E.B., and Webster, A.R. 1935.

Churchill Mining and Milling Company, Limited; p.82 *in* Mines of Ontario in 1934; Ontario Dept. Mines Vol.44, pt.I, p.67-167 (published 1936).

Sinclair, D.G., Reeley, B.C., Cooper, D.F., Weir, E.B., and Webster, A.R. 1936.

Churchill Mining and Milling Company, Limited; p.92 *in* Mines of Ontario in 1935, Ontario Dept. Mines, Vol.45, pt.I, p.77-188 (published 1937).

Terraquest Ltd., and MacGregor, R 2008.

Operations Report for Platinex Inc. High Resolution Tri-Sensor Magnetic, XDS VLF-EM and Radiometric Airborne Survey.

Tindale, J. L. 1989.

Summary Report of Work Performed on the Shining Tree Property, Asquith Township During 1988-1989 Season of Asquith Resources Inc. Assessment File 41P11SE0152.

Tomlinson, K.Y., Thurston, P.C., Hughes, D.J. and Keays R.R. 1996c.

The central Wabigoon region: Petrogenesis of mafic-ultramafic rocks in the Steep Rock, Lumby Lake and Obonga Lake greenstone belts (continental drifting and rifting in the Archaean); Harrap R.M. and Helmstaedt H. (eds)., Lithoprobe Secretariat, University of British Columbia, Lithoprobe Report 53, p.65-73.

Trip, G. 2000.

Structural Geology and Gold Mineralisation of the Ora Banda and Zuleika districts, Eastern Goldfields, Western Australia, Master of Science thesis abstract.

Trusler, I and Trusler, J 2012 – Assessment Report on Diamond Drilling of the Herrick Gold Deposit. Churchill, Larder Lake Mining Division, Ontario, Canada.

Xie, Q. and Kerrich, R. 1994.


Silicate-perovskite and majorite signature komatiites from the Archaean Abitibi Greenstone Belt: Implications for early mantle differentiation and stratification; Journal of Geophysical Research, v.99, p.15799-15812.

25.0 CERTIFICATES OF QUALIFICATIONS

I, James R. Trusler at 11 Algonquin Crescent, Aurora, Ontario do hereby certify that:

- 1) I am a Geological Engineer employed as President and CEO of Platinex Inc. and I am also the principal shareholder of Platinex Inc.;
- 2) I graduated from the University of Toronto with BA.Sc. in Geological Engineering in 1967. I obtained a Master of Science (Geology) from Michigan Technological University in 1972. I have practiced my profession full-time from 1967-1969 and from 1970 to present;
- 3) I am a Professional Engineer registered with the Professional Engineers Ontario (PEO #47064019);
- 4) I have not received, nor do I expect to receive, any interest, directly or indirectly, in the Shining Tree property;
- 5) As of the date of this certificate, to the best of my knowledge, information and belief, this report contains all scientific and technical information that is required to be disclosed to make the Summary Report on the Shining Tree Gold property not misleading;
- 6) I have read National Instrument 43-101 and written the Summary Report on the Shining Tree Gold property in compliance with the intent of National Instrument 43-101 and Form 43-101F1 but is not a Technical Report as defined by National Instrument 43-101;
- 7) I have relied on and quoted extensively from the 2008 Qualifying Report prepared by Garry Bryant and David R. Jamieson;
- 8) I have relied on discussions and records of exploration recorded by and supervised by David R. Jamieson, Scott Franko and Dean Cutting all of whom are independent qualified persons;
- 9) I have visited the property on several occasions during the exploration programs from 2008 to 2012.

Dated the 18th Day of June, 2012

A circular seal for a Registered Professional Engineer in the Province of Ontario. The seal contains the text "REGISTERED PROFESSIONAL ENGINEER" and "PROVINCE OF ONTARIO". A signature is written across the seal, and the name "J. R. TRUSLER" is printed below it.

Signature of Qualified Person:

James R. Trusler, B.A.Sc, MS, P.Eng

Summary Report for Platinox Inc., Shining Tree, Ontario

By: James R. Trusler

TABLE 1					
Shining Tree Project - Claims and Status as of June 7, 2012					
Township/Area	Claim Number	Units	Recording Date	Claim Due Date	Total Reserve
ASQUITH	4203475	2	2005-Nov-29	2013-Nov-29	\$0
ASQUITH	3011745	3	2004-Aug-17	2013-Aug-17	\$0
ASQUITH	4201206	4	2005-Nov-29	2013-Nov-29	\$0
ASQUITH	4207969	4	2005-May-30	2014-May-30	\$0
ASQUITH	4209215	4	2006-Jan-03	2014-Jan-03	\$0
ASQUITH	4211939	3	2006-Nov-20	2013-Nov-20	\$0
ASQUITH	4217643	9	2007-Jan-19	2014-Jan-19	\$0
CHURCHILL	1192177	2	2002-Jul-22	2014-Jul-22	\$0
CHURCHILL	1199655	2	2002-May-14	2014-May-14	\$0
CHURCHILL	1217520	1	1996-Nov-25	2013-Nov-25	\$0
CHURCHILL	1217521	1	1996-Nov-25	2013-Nov-25	\$0
CHURCHILL	1225095	1	1997-Jun-18	2013-Jun-18	\$0
CHURCHILL	1226938	1	1998-Jun-15	2014-Jun-15	\$0
CHURCHILL	1227175	1	1998-Sep-29	2013-Sep-29	\$0
CHURCHILL	1235004	1	2000-Aug-16	2013-Aug-16	\$278
CHURCHILL	1235157	2	2000-Jun-13	2013-Jun-13	\$0
CHURCHILL	1238874	1	1999-Jun-18	2014-Jun-18	\$0
CHURCHILL	1238875	1	1999-Jun-17	2013-Jun-17	\$0
CHURCHILL	1238881	2	1999-Jun-21	2013-Jun-21	\$0
CHURCHILL	1242019	1	2000-Jun-22	2016-Jun-22	\$703,448
CHURCHILL	1242194	1	2001-Apr-10	2014-Apr-10	\$0
CHURCHILL	1242933	1	2000-Nov-29	2013-Nov-29	\$0
CHURCHILL	1242934	1	2000-Nov-27	2013-Nov-27	\$0
CHURCHILL	3004540	1	2002-Sep-12	2013-Sep-12	\$0
CHURCHILL	3011224	2	2003-Apr-28	2014-Apr-28	\$0
CHURCHILL	3011227	3	2003-Jul-23	2014-Jul-23	\$0
CHURCHILL	3011743	4	2004-Aug-17	2013-Aug-17	\$0
CHURCHILL	3014015	8	2005-Jan-05	2014-Jan-05	\$393
CHURCHILL	3014016	1	2005-Jan-05	2014-Jan-05	\$0
CHURCHILL	4203474	4	2005-Jan-05	2014-Jan-05	\$0
CHURCHILL	4207970	2	2005-May-30	2014-May-30	\$0
CHURCHILL	4207971	3	2005-May-30	2014-May-30	\$0
CHURCHILL	4207972	2	2005-May-30	2013-May-30	\$1,423
CHURCHILL	4209217	1	2006-Jan-03	2014-Jan-03	\$0
CHURCHILL	4209231	3	2008-Dec-03	2013-Dec-03	\$0
CHURCHILL	4211938	10	2006-Nov-20	2013-Nov-20	\$0
CHURCHILL	4217645	6	2007-Jan-19	2013-Jan-19	\$0
CHURCHILL	4217646	12	2007-Jan-19	2014-Jan-19	\$0
CHURCHILL	4245864	1	2008-Dec-03	2013-Dec-03	\$0
MACMURCHY	1192173	1	2002-Oct-29	2013-Oct-29	\$0
MACMURCHY	4201272	3	2007-Aug-24	2013-Aug-24	\$0
MACMURCHY	4203531	3	2005-Feb-24	2013-Feb-24	\$0
MACMURCHY	4207973	2	2005-May-30	2014-May-30	\$0
MACMURCHY	4207974	3	2005-May-30	2014-May-30	\$0
MACMURCHY	4217644	12	2007-Jan-19	2014-Jan-19	\$0
MACMURCHY	4252127	3	2009-Dec-07	2017-Dec-07	\$1,298
TOTAL		139			\$706,840

Summary Report for Platinex Inc., Shining Tree, Ontario

By: James R. Trusler

Note 1: It is understood from Iain Trusler that, assessment submittals to date reserve on file and available additional expenditures will keep all claims in good standing well into the future and possibly in excess of twenty years.

Note 2: A review of the Mining Records published records confirmed that all claims are valid and the owner of record is Platinex Inc.

APPENDIX A: TABLE 2 GLOSSARY AND ABBREVIATIO OF TERMS

Term or Abbreviation	Meaning
AEM	Airborne Electromagnetic
Ag	silver
Al	Aluminum
AMSL	above mean sea level
Au	gold
Be	beryllium
BGC	Bryant Groundwater Consulting
Cd	cadmium
Cu	copper
EM	electromagnetic
FA/ICP-MS	Fire assay/inductively coupled plasma-mass spectrometer
g	gram(s)
g/t	grams per tonne (equivalent to ppm)
GPS	Global Positioning Systems
GSC	Geological Survey of Canada
Ha	Hectare(s) (2.471 acres)
kg	kilogram(s)
km	kilometre(s)
KM	Kidd-Munro
m	metres
MNDM	Ministry of Northern Development and Mines
MSL	Mean Sea Level (0 m)
Ni	Nickel
ODM	Overburden Drilling Management
OGS	Ontario Geological Survey
oz	Troy ounce (31.1035 grams)
oz/t	Troy ounces per tonne
Pb	lead
Pd	palladium
ppm, ppb	parts per million/parts per billion
Pt	platinum
REE	Rare Earth Elements
REE	rare-earth elements
Ti	Titanium
tonnes or t	metric tones


APPENDIX B: CERTIFICATE OF CONSENT

To accompany the report entitled as above

I, James R. Trusler, residing at 11 Algonquin Crescent, Aurora, Ontario do hereby certify that:

- 1) I am an Geological Engineer but not independent of Platinex;
- 2) I graduated from the University of Toronto with a Bachelor of Applied Science BASc 1967. I obtained a Master of Science (Geology) from Michigan Technological University in 1972. I have practiced my profession full-time from 1967-1969 and from 1970 to present;
- 3) I am a Professional Engineer registered with the Professional Engineers Ontario of the province of Ontario (PEO #47064019);
- 4) I have not received, nor do I expect to receive, any interest, directly or indirectly, in the Shining Tree project but am a major shareholder, President, CEO and director of Platinex Inc.;
- 5) As of the date of this certificate, to the best of my knowledge, information and belief, this technical report contains all scientific and technical information that is required to be disclosed to make the summary report not misleading;
- 6) I have read National Instrument 43-101 and this summary report has been prepared in compliance with National Instrument 43-101 and Form 43-101F1;
- 7) I am not independent of the issuer as defined in Section 1.4 of National Instrument 43-101;
- 8) I am the sole author of this report;
- 9) I visited the property on numerous occasions during the exploration and became familiar with all project areas. I also examined showings, and chip and core samples for validation of gold mineralization grades were collected for the 2008 Qualifying Report by Bryant and Jamieson;
- 10) I prepare the summary report for the Shining Tree Gold Property in accordance with NI 43-101 and Form 43-101 FI guidelines. The report is based on the review of published information, company files and discussions with Platinex Inc. Personnel and site visits;
- 11) I consent to the filing of the Summary Report with any stock exchange and other regulatory authority and any publication by them, including electronic publication. I also consent to Platinex Inc. posting of this Summary Report on their corporate website which is accessible to the public.

Signed



James R. Trusler



Dated: June 18, 2012

APPENDIX C: TABLE D1 – D4: HISTORIC CHECK ASSAY AND PULP METALLIC ASSAYS PERFORMED ON HERRICK DEPOSIT SAMPLES

TABLE D1: UNOCAL 1989 CHANNEL SAMPLE GOLD ASSAYS

<i>Sample No</i>	1st PULP		2nd PULP		3rd PULP	
	<i>Au Assay (g/t)</i>	<i>Check Assay (g/t)</i>	<i>2nd Au Assay (g/t)</i>	<i>Check Au Assay (g/t)</i>	<i>3rd Au Assay (g/t)</i>	<i>Check Au Assay (g/t)</i>
6519	34.35	26.57	25.41	22.22		
7481	23.18	27.57	25.54	26.64		
6052	7.44	7.27	7.1	6.72		
6063	18.51	16.83	19.27	19.2		
6093	9.6	8.74	11.31	9.15		
6126	61.03	67.2	33.43	31.2	73.71	72.69
7604	11.73	11.9	11.73	12.21		

Note: All assays were performed by Swastika Laboratories which was not an ISO accredited laboratory in 1989-1990. Documents include copies of Geochemical Analysis Certificates.

TABLE D2: UNOCAL 1989 SPLIT DRILL CORE SAMPLES GOLD ASSAYS

<i>Sample No</i>	1ST PULP		2ND PULP		
	<i>Au Assay (g/t)</i>	<i>Check Assay (g/t)</i>	<i>2nd Assay (g/t)</i>	<i>Check Assay (g/t)</i>	<i>Pulp metallic (g/t)</i>
14521	4.66	4.87	4.25	4.29	
14608	6.34	8.3	8.47	8.54	
14654	17.14	15.12	15.81	16.73	
14684	15.5	16.46	17.21	15.67	
14713	8.47	8.81	7.47	7.75	8.09
14815	6.24	6.62	6.93	6.86	7.08
14922	6.48	7.06	9.36	10.01	8.25
14955	4.25	5.66			5.96
14950	6.38	6.31	7.1	6.41	5.56
14875	2.3	2.33			2.87
14865	7.47	7.03	6.58	7.37	7.87

Note: All assays were performed by Swastika Laboratories which was not ISO accredited laboratory in 1989. Documents include copies of Geochemical Analysis Certificates.

TABLE D3: UNOCAL 1989 PULP AND METALLIC ASSAYS

Sample No	Au Assay (g/t) +100 mesh	Au Assay (g/t) - 100 mesh	Au Assay (g/t) - 100 mesh	Calculated Au value (g/t)
14875	8.08	2.91	2.81	2.87
14914	9.65	5.49	5.42	5.56
14922	5.67	7.95	8.57	8.25
14950	4.71	5.62	5.52	5.56
14713	16.32	7.1	8.81	8.09

Note: These five samples had the highest concentrations of gold in the metallic fraction. A total of 125 samples were submitted by Unocal for pulp and metallic assays. Swastika Laboratories performed the analysis and was not ISO certified. Documents include copies of Geochemical Analysis Certificates.

TABLE D4: FORT KNOX 1990 CHANNEL SAMPLE GOLD ASSAYS

Sample No	1ST PULP		2ND PULP	
	(ppb Au)	Check Assay (ppb Au)	2 nd Assay (ppb Au)	2nd Pulp Check Assay (ppb Au)
122835	8,297	6,994		
122846	24,754	27,566		
122847	22,766	22,560	30,515	28,252
122869	13,714	12,412		
122870	18,103	19,543	17,966	19,474

Note: All assays were performed by Swastika Laboratories which was not an ISO accredited laboratory in 1989-1990. Documents include copies of Geochemical Analysis Certificates.



500 Km



Source: National Topographic Data Base - GeoGratis
OGS MRD 126

PLATINEX Shining Tree Project
The Quest for a Greener Planet

GENERAL LOCATION MAP

Drawn by: IST


Checked by: JRT

Scale: 1:9,000,000

Date: June 2012

I.D.:STloc9m040612

FIGURE 1



Source: National Topographic Data Base - GeoGratis

NTS 41P11


Symbol

- Shining Tree Property
- Creso Exploration Claims
- Townships
- Joint Venture Boundary
- Primary Roads
- Bush Roads and Former Roads

PLATINEX The Quest for a Greener Planet **Shining Tree Project**

CLAIM MAP

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STcl50k040612	FIGURE 2



Shining Tree Project <small>The Quest for a Greener Planet</small>		
GEOLOGICAL SUBDIVISIONS OF ONTARIO		
Drawn by: IST	Checked by: JRT	Scale: 1:9,000,000
Date: June 2012	I.D.:STsub9m070612	FIGURE 3





Figure 4

Simplified Geological Map of the Abitibi Greenstone Belt Showing Distribution of Major Fault Zones and Gold Deposits (after Dube and Gosselin, 2005; Poulsen et al, 2000).  Platinox Shining Tree Gold Property





Figure 5: **Lithostratigraphic Assemblages of the Shining Tree Area**
 (after Oliver et al, 1999b, Figure 6.1)

June 2012


- | | | |
|-----------------------------------|----------------------------------|------------------------------------|
| Proterozoic Rocks | Intrusive contact | Tisdale Assemblage (2,707.5 Ma) |
| Timiskaming Age Intrusive Rocks | Intrusive contact / unconformity | Kidd-Munro Assemblage (2,716.7 Ma) |
| Timiskaming Assemblage (2,687 Ma) | Intrusive contact | Deloro Assemblage (2,726.5 Ma) |
| Keewatin Intrusive Rocks | Unconformity | Pacaud Assemblage (2,741 Ma) |
| Intrusive contact | Fawcett igneous structure | Fault |
| | | Interpreted contact |
| | | Assemblage boundary |













PLATINEX The Quest for a Greener Planet **Shining Tree Project**


GEOLOGY

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STgrl50k040612	FIGURE 6



Source: M.Spaho, after G.R.Cluff 1989.


Symbol	
	Diabase
	Trachyte
	Greywacke
	Polymictic Conglomerate
	Volcanic Breccia
	Lapilli Tuff
	Mineralized Zone (At Surface)
	Mineralized Zone (In Drill Holes)
	Herrick Shaft
	Diamond Drill Holes



Shining Tree Project

HERRICK GEOLOGY AND DRILL HOLE LOCATIONS

Drawn by: IST	Checked by: JRT	Scale: 1:2,000
Date: June 2012	I.D.:STher2k060612	FIGURE 7



Map objects scale and location are approximate


Symbol

- Veins (1-17/101-123)
- - - Inferred Veins
- Caswell Shafts
- Silver (Ag 0.1 - 2.15 g/t)
- Tellurium (Te 0.1 - 1.1ppm)
- Chesbar Grab Samples c1987 (Au g/t)
- Channel Samples (Au g/t) West side 2008, East side 2010
- Drill Holes
- 2011 DDH (WP11)

PLATINEX Shining Tree Project
The Quest for a Greener Planet

CASWELL PROSPECT HIGHLIGHTS

Drawn by: IST	Checked by: JRT	Scale: 1:3,500
Date: June 2012	I.D.:STcas3.5k070612	FIGURE 8







BP12-02
Depth 200m
EOH

175
150
125
100
75
50

20 Metres

NTS 41P11

Symbol

-  Pillowed Mafic Volcanics
-  Felsic Volcaniclastics
-  Gabbro
-  Channel Sample

PLATINEX Shining Tree Project
The Quest for a Greener Planet

BEILBY LAKE SHOWING

Drawn by: IST


Checked by: JRT

Scale: 1:500

Date: June 2012

I.D.:STbil5h070612

FIGURE 9




- Symbol**
- Shining Tree Property
 - Townships
 - Primary Roads
 - Bush Roads and Former Roads

PLATINEX Shining Tree Project
The Quest for a Greener Planet

**PRELIMINARY IP SURVEY PLANS
CHARGEABILITY N=4**


Drawn by: IST	Checked by: JRT	Scale: 1:20,000
Date: June 2012	I.D.:STchg20k070612	FIGURE 10



Source: OGS Geophysical Data Sets 1003, 1064.


Symbol	Electromagnetic Decay Constant Classification
	● > 3,000 microseconds
	● 2,000 to 3,000
	● 1,000 to 2,000
	● 500 to 1,000
	⊗ 250 to 500
	○ 0 to 250

<h2>Shining Tree Project</h2>		
<h3>FIRST VERTICAL DERIVATIVE MAGNETICS AND EM ANOMALIES</h3>		
Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STma1d50k060612	FIGURE 11



Source: OGS Geophysical Data Sets 1003.


Symbol	Electromagnetic Decay Constant Classification
	● > 3,000 microseconds
	● 2,000 to 3,000
	● 1,000 to 2,000
	● 500 to 1,000
	● 250 to 500
	○ 0 to 250



Shining Tree Project

XDS ORTHO AND EM ANOMALIES

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.: STort50k060612	FIGURE 12




Source: Geophysical Interpretation by Z. Dvorak, Independent Geophysical Consultant

- Shining Tree Property
- Townships
- Primary Roads
- Dykes
- A. Magnetic low outline of intrusive
- B. Triangular magnetic high area
- VO1 - VO4 VLF-EM ortho anomalies
- VL1 - VL5 VLF-EM line anomalies
- Faults or fractures interpreted from magnetic pattern
- M1 - M9 Magnetic anomalies
- RK1 - RK5 Radiometric (potassium) anomalies

PLATINEX Shining Tree Project
The Quest for a Greener Planet

**INTERPRETATION OF
 GEOPHYSICAL SURVEY RESULTS**

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STint50k060612	FIGURE 13







FIGURE: 15


Relation of Gold in Till to Major Gold Deposits

Shining Tree Project

Source: MNDMF GDS 1036, Various Reports

 Gold Mine/Deposit
 Gold Grains In Till

 200 and up
 100
 20

Gravity
 mGal
 High 
 Low

Cut off: 10 gold grains per 10kg sample
 Based on 3988 till samples within 66,000sq km
 (NTS 41O, 41P, 42A, 42B)






Figure: 16



NTS 41R11


- Symbol**
- Shining Tree Property
 - Glacial Gold Trains
 - Primary Roads
 - Bush Roads and Former Roads
 - Historic Mines/Shafts
 - Historic Gold Showings



PLATINEX The Quest for a Greener Planet **Shining Tree Project**


GOLD GRAINS IN TILL

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STtill50k070612	FIGURE 17



- PROTEROZOIC**
- 20 Matachewan Diabase
 - 19 Nippissing Gabbro
- ARCHEAN**
- 14 Intermediate to Felsic Intrusive Porphyry
 - 10 Clastic Metasedimentary Rocks
 - 8 Intermediate to Felsic Intrusive Rocks
 - 7 Intermediate to Felsic Meta-Porphyry Intrusives
 - 6 Mafic to Ultramafic Meta-Intrusive Rocks
 - 4 Chemical Metasedimentary Rocks
 - 3 Intermediate to Felsic Metavolcanic Rocks
 - 2 Mafic to Intermediate Metavolcanic Rocks
 - 1 Ultramafic Metavolcanic Rocks


- Symbol**
- Shining Tree Property
 - XDS Ortho Anomalies
 - Deformation Zone (Johns, 1996)
 - Bush Roads and Former Roads
 - Historic Mines/Shafts
 - Historic Gold Showings



PLATINEX The Quest for a Greener Planet **Shining Tree Project**


COMPILATION: GEOLOGY, XDS ORTHO ANOMALIES WITH GOLD IN TILL

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STcom50k070612	FIGURE 18



NTS 41R11


- Symbol**
- Shining Tree Property
 - Townships
 - Primary Roads
 - Bush Roads and Former Roads
 - Historic Mines/Shafter
 - Historic Gold Showings



PLATINEX The Quest for a Greener Planet **Shining Tree Project**


ARSENIC IN TILL

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STas50k070612	FIGURE 19



NTS 41R11


- Symbol**
- Shining Tree Property
 - Townships
 - Primary Roads
 - Bush Roads and Former Roads
 - Historic Mines/Shafter
 - Historic Gold Showings



PLATINEX The Quest for a Greener Planet **Shining Tree Project**


NICKEL IN TILL

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STni50k070612	FIGURE 20



NTS 41P11


- Symbol**
- Shining Tree Property
 - Townships
 - Primary Roads
 - Bush Roads and Former Roads
 - Historic Mines/Shafts
 - Historic Gold Showings



PLATINEX The Quest for a Greener Planet **Shining Tree Project**


COPPER IN TILL

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.:STcu50k070612	FIGURE 21



NTS 41P11

- Symbol**
- Shining Tree Property
 - Townships
 - Primary Roads
 - Bush Roads and Former Roads
 - Historic Mines/Shafts
 - Historic Gold Showings



PLATINEX The Quest for a Greener Planet **Shining Tree Project**

SULPHUR IN TILL

Drawn by: IST	Checked by: JRT	Scale: 1:50,000
Date: June 2012	I.D.: STs50k070612	FIGURE 22